

CHEMISTRY NOTES FOR CLASS - 11 (FOR PUNJAB)

Chapter

09

ACID-BASE CHEMISTRY

Student Learning Outcomes

After studying this chapter, students will be able to:

- Define conjugate acid base pairs. (Understanding)
- Identify conjugate acid base pairs in reactions. (Understanding)
- Distinguish that Lewis acids accept lone pair, and Lewis bases donate lone pair to make a coordinate covalent bond. (Understanding)
- Define mathematically the terms K_w , pH, K_a and pK_a and use them in calculations. (K_b and the equation $K_w = K_a \times K_b$ will not be tested). (Application)
- Calculate $H^+_{(aq)}$ and pH values for: (Application)
 - (a) strong acids (b) strong alkalis (c) weak acids (d) weak alkalies.
- Calculate the $[H_3O^+]$ given the K_a and molar concentration of weak acid. (Application)
- Apply the concept of the common ion effect to describe why the solubility of a substance changes when it is dissolved in a solution containing a common ion. (Application)
- Calculate the pH of buffer solutions in given appropriate data. (Application)
- Construct an expression for K_{sp} . (Application)
- Demonstrate the ability to comprehend and effectively apply the concept of solubility product K_{sp} . (Application)
- Perform calculations using K_{sp} values and concentration of a common ion. (Application)
- Calculate concentrations of ions of slightly soluble salts. (Application)
- Calculate K_{sp} from concentrations and vice versa. (Application)
- Use the concept of hydrolysis to explain why aqueous solutions of some salts are acidic or basic. (Understanding)
- Apply the concept of conjugate acid and conjugate base on salt hydrolysis. (Application)
- Select suitable indicators for acid-alkali titration, given appropriate data (pK_a values will not be used). (Understanding)
- Perform acid-base titration to calculate molarity and strength of given sample solutions. (Application)

Differences between Common Acids and Bases

Sr.No.	Acids	Bases
1.	Acids were first recognized as having sour taste.	Bases were first recognized as having bitter taste.
2.	Acids change the litmus from blue to red.	Bases change the litmus from red to blue.
3.	Acids change the indicator "Phenolphthalein" from pink to colourless.	Bases change the indicator "Phenolphthalein" from colourless to pink.
Note: Acids and bases neutralize each other and during neutralization, acids and bases react with each other to produce ionic substances called salts.		



Interesting Information!

Acidity in our stomachs is due to excess HCl. It is treated by taking mild bases such as, baking soda.

CHEMISTRY NOTES FOR CLASS - 11 (FOR PUNJAB)

BRONSTED-LOWRY CONCEPT

[Exercise L.Q.3]

Q. Describe the Bronsted-Lowry theory of acids and bases. Provide examples of conjugate acid-base pairs and explain clearly their relationship.

The Danish chemist J.N. Bronsted and the English chemist T.M. Lowry in 1923 independently expanded the Arrhenius theory of acid-base.

➤ **Bronsted-Lowry Acid:**

According to the Bronsted-Lowry concept, "an acid is the species donating a proton in a proton-transfer reaction".


➤ **Bronsted-Lowry Base:**

According to Bronsted-Lowry concept, "a base is the species accepting the proton".

Example:

When HCl dissolves in Water:

When HCl dissolves in water, an H^+ ion (a proton) is transferred from HCl to water, where it becomes attached to a lone pair of electrons on the O atom and forms H_3O^+ . In fact, HCl (the acid) has donated the H^+ and H_2O (the base) has accepted it.



➤ **Conjugate Acid:**

"A species formed after a Bronsted base accepts a proton from the acid is called the conjugate acid". The hydronium ion (H_3O^+) is the conjugate acid of water.


➤ **Conjugate Base:**

"A species formed when an acid donates a proton to a base is called the conjugate base". $HCl-Cl^-$ and $H_2O-H_3O^+$ are conjugate acid-base pairs.

Example:

When NH_3 dissolves in Water:

When ammonia dissolves in water, proton transfer also occurs. An H^+ from H_2O attaches to the N atom's lone pair and NH_4^+ and OH^- are formed. With fewer H^+ , the water molecule becomes OH^- ion:



In this case, H_2O (the acid) has donated the proton and NH_3 (the base) has accepted it. $NH_3-NH_4^+$ and H_2O-OH^- are conjugate acid-base pair.

Amphoteric Substance: An amphoteric substance is a species that can act as either an acid and a base (it can lose or gain a proton), depending on the other reactant.

H_2O is Amphoteric in Nature: H_2O is amphoteric in nature because it acts as a base in one case and as an acid in the other (as shown in the above examples).



The limitations of Arrhenius theory is that it describes the reaction of an acid and base only in the aqueous medium. There are many reactions that occur in solvents other than water or in the absence of any solvent. Bronsted-Lowry pointed out that acid-base reactions can be seen as proton transfer reactions and that acids and bases can be defined in terms of this proton (H^+) transfer.

Note:

- Chloride ion (Cl^-) is indeed the conjugate base of hydrochloric acid (HCl) because it is the species that remains after the acid (HCl) has lost its proton.

- A Bronsted-Lowry acid is a proton donor.
- A Bronsted-Lowry base is a proton acceptor.



Rack Your Mind!

- In the reaction between HCl and water, which species acts as the conjugate base?
A) H_3O^+ B) Cl^-
C) HCl D) H_2O




Rack Your Mind!

- Conjugate acid-base pair differs by:
A) An electron pair B) A neutron
C) An electron D) A proton

CHEMISTRY NOTES FOR CLASS - 11 (FOR PUNJAB)

Application of Bronsted-Lowry Concept: The Bronsted-Lowry theory can be applied to acids in solvents other than water or even solventless reactions. The reaction between gaseous ammonia and HCl gives solid NH_4Cl .



HCl is an acid because in the reaction it donates a proton to the NH_3 molecule. The NH_3 acts as a base, even though hydroxide ion OH^- is not present, and accepts a proton from the HCl molecule. Similarly,




Table: Conjugate Acid-Base Pairs of Common Species

Acid	Base	Conjugate Acid	Conjugate Base
HNO_3	$+$ H_2O	$\rightleftharpoons$ H_3O^+	$+$ NO_3^-
H_2SO_4	$+$ H_2O	$\rightleftharpoons$ H_3O^+	$+$ HSO_4^-
H_2O	$+$ CO_3^{2-}	$\rightleftharpoons$ HCO_3^-	$+$ OH^-

LEWIS CONCEPT OF ACIDS AND BASES

(Exercise L.Q.4)

Q. Define the Lewis theory of acids and bases. How does this theory differ from the Bronsted-Lowry theory? Give examples of Lewis acids and bases that do not involve proton transfer.

In 1923, G.N. Lewis proposed a generalized definition of acid-base behavior in which acids and bases are identified by their ability to accept or to donate a pair of electrons and form a coordinate covalent bond.

➤ **Lewis Acid:**

"A Lewis acid is any species (molecule or ion) that can accept a pair of electrons".


➤ **Lewis Base:**

"A Lewis base is any species (molecule or ion) that can donate a pair of electrons".

A Lewis acid-base reaction occurs when a base donates a pair of electrons to an acid.

Example 1:

In the following reaction, each of two ammonia molecules, Lewis bases, donates a pair of electrons to a positively charged silver ion, the Lewis acid. The sum of charges on the left side is +1, so the acid-base adduct on the right hand side must carry a charge of +1:



Example 2:

The boron atom in boron trifluoride, BF_3 , has only six electrons in its valence shell. Boron atom has an incomplete octet. So it can behave as an electron pair acceptor. BF_3 is a very good Lewis acid and reacts with many Lewis bases e.g. a fluoride ion which is the Lewis base in this reaction, donates one of its lone pairs:




CHEMISTRY NOTES FOR CLASS - 11 (FOR PUNJAB)

The negative charge on the adduct is the sum of charges on the left hand side of the equation is -1 , the sum of charges on the right hand side must also be -1 .


In the following reaction NH_3 has a lone pair on N-atom so it is a Lewis base (electron pair donor) and boron accepts the lone pair and is a Lewis acid.

Example 3:



QUICK CHECK 9.1

a) Identify the conjugate acid-base pairs in the following reactions:



Also write down the balanced chemical equation for the reaction and explain.


Ans. The conjugate acid-base pairs are:

1. HCN (acid) and CN^- (its conjugate base)
 HCN donates a proton (H^+) to become CN^- .
2. H_2O (base) and H_3O^+ (its conjugate acid)
 H_2O accepts a proton (H^+) to become H_3O^+ .

Summary of Pairs:

Acid	Conjugate Base
HCN	CN^-
H_2O	H_3O^+

- Acid $\rightarrow$ Conjugate base by losing H^+ .
- Base $\rightarrow$ Conjugate acid by gaining H^+ .



Ans. The conjugate acid-base pairs are identified as follows:

Pair 1: CH_3COOH (Acid) and CH_3COO^- (Conjugate Base)

- CH_3COOH donates a proton (H^+) to become CH_3COO^- .
 - Acid: CH_3COOH (acetic acid).
 - Conjugate Base: CH_3COO^- (acetate ion).

Pair 2: H_2O (Base) and H_3O^+ (Conjugate Acid)

- H_2O accepts a proton (H^+) to become H_3O^+ .
 - Base: H_2O (water).
 - Conjugate Acid: H_3O^+ (hydronium ion).



Ans. Conjugate Acid-Base Pairs:


- H_2S (acid) / HS^- (Conjugate base): H_2S donates a proton (H^+) to become HS^- . (Acid: Proton (H^+) donor.)
- H_2O (base) / H_3O^+ (Conjugate acid): H_2O accepts a proton (H^+) to become H_3O^+ . (Base: Proton (H^+) acceptor.)
- Pairs differ by one proton (H^+).

Example:

- H_2S (acid) $\rightarrow$ HS^- (conjugate base)
- H_2O (base) $\rightarrow$ H_3O^+ (conjugate acid)

Note: H_3O^+ is the strongest acid here; HS^- is the weakest base.

CHEMISTRY NOTES FOR CLASS - 11 (FOR PUNJAB)



Ans. Conjugate Acid-Base Pairs:

1. HCl/Cl^-

- HCl (acid) donates a proton (H^+) to become Cl^- (its conjugate base).

2. $\text{HCO}_3^-/\text{H}_2\text{CO}_3$

- HCO_3^- (base) accepts a proton (H^+) to become H_2CO_3 (its conjugate acid).

Summary Table:

Acid	Conjugate Base
HCl	Cl^-
H_2CO_3	HCO_3^-

- Acids lose H^+ to form conjugate bases.

- Bases gain H^+ to form conjugate acids.

b) Identify the Lewis acid and Lewis base in the reaction between:

i. SO_3 and O^{2-}

Ans. Balanced Chemical Equation:



Lewis Acid-Base Identification:

1. Lewis Acid: SO_3

- Reason: SO_3 accepts an electron pair from O^{2-} (its sulfur atom has an incomplete octet and is electron-deficient).

2. Lewis Base: O^{2-}

- Reason: O^{2-} donates an electron pair to SO_3 (it has lone pairs available for bonding).

Mechanism Explanation:

1. SO_3 (electron-deficient due to polar $\text{S}=\text{O}$ bonds) acts as the electron-pair acceptor (Lewis acid).

2. O^{2-} (a negatively charged ion with lone pairs) acts as the electron-pair donor (Lewis base).

3. The O^{2-} attacks the sulfur in SO_3 , forming a coordinate covalent bond and creating SO_4^{2-} (sulfate).


Key Notes:

- SO_3 is a classic Lewis acid (e.g., in sulfuric acid production: $\text{SO}_3 + \text{H}_2\text{O} \longrightarrow \text{H}_2\text{SO}_4$).

- O^{2-} is a strong Lewis base (common in oxide reactions).

ii. HCl and H_2O

Ans. Balanced Equation:



Lewis Acid-Base Identification:

1. Lewis Acid: H^+ (from HCl)

- Reason: Accepts an electron pair from H_2O to form H_3O^+ .

2. Lewis Base: H_2O

- Reason: Donates an electron pair to H^+ to form a coordinate covalent bond in H_3O^+ .

Explanation:

- HCl dissociates into H^+ (electron-pair acceptor, Lewis acid) and Cl^- .

- H_2O (with lone pairs on oxygen) acts as the electron-pair donor (Lewis base), forming hydronium (H_3O^+).

iii. BF_3 and NH_3

Ans. Balanced Chemical Equation:



Lewis Acid and Base Identification:

- Lewis Acid: BF_3 (electron pair acceptor).

- Boron in BF_3 has an empty $2p$ orbital, making it electron-deficient.

CHEMISTRY NOTES FOR CLASS - 11 (FOR PUNJAB)

- **Lewis Base:** NH_3 (electron pair donor).
 - Nitrogen in NH_3 has a **lone pair** it can donate.

Mechanism:

1. NH_3 donates its lone pair to the empty orbital of BF_3 .
2. A **dative covalent bond** (coordinate bond) forms between N and B.


Key Points:

- BF_3 is a classic Lewis acid (no protons involved!).
- NH_3 is a Lewis base due to its lone pair.
- The product ($\text{H}_3\text{N}-\text{BF}_3$) is a stable adduct.

IONIC PRODUCT OF WATER

- Pure water is a poor conductor of electricity but its conductance is measurable.
- Water undergoes self-ionization reversibly as follows,

Net Reaction:



The equilibrium constant for this reaction can be written as follows:

$$K_c = \frac{[\text{H}^+][\text{OH}^-]}{[\text{H}_2\text{O}]} = 1.8 \times 10^{-16} \text{ mol dm}^{-3}$$

- The concentration of H_2O i.e. $[\text{H}_2\text{O}]$ in pure water may be calculated to be 1000 g dm^{-3} divided by 18 g mol^{-1} giving $55.5 \text{ moles dm}^{-3}$ i.e. $\frac{1000 \text{ g dm}^{-3}}{18 \text{ g mol}^{-1}} = 55.5 \text{ mol dm}^{-3}$
- As water is present in very large excess and very few of its molecules undergo ionization, so its concentration remains effectively constant.
- Constant concentration of water is taken on L.H.S. and multiplied with K_c to get another constant called K_w .

$$K_c[\text{H}_2\text{O}] = [\text{H}^+][\text{OH}^-]$$

$$\text{or} \quad = 1.8 \times 10^{-16} \times 55.5 = 1.01 \times 10^{-14} = [\text{H}^+][\text{OH}^-]$$

This 1.01×10^{-14} is called K_w of water at 25°C .

$$K_c[\text{H}_2\text{O}] = [\text{H}^+][\text{OH}^-]$$

So, $K_w[\text{H}_2\text{O}] = [\text{H}^+][\text{OH}^-] = 10^{-14}$ at 25°C .

- K_w is called ionic product of water or dissociation constant of water.

➤ K_w of H_2O Increases with Increase of Temperature:

- The value of K_w increases almost 75 times when temperature is increased from 0°C to 100°C .

For neutral water

$$[\text{H}^+] = [\text{OH}^-]$$

Or

$$[\text{H}^+][\text{H}^+] = 10^{-14}$$

$$[\text{H}^+]^2 = 10^{-14} \text{ (at } 25^\circ\text{C)}$$

$$[\text{H}^+] = 10^{-7} \text{ mol dm}^{-3}$$

And

$$[\text{OH}^-] = 10^{-7} \text{ mol dm}^{-3}$$

The effect of temperature on K_w is shown in Table.

- Whenever some quantity of acid or base is added to water, then K_w remains the same, but $[\text{H}^+]$ and $[\text{OH}^-]$ are no more equal.



Rack Your Mind!

4. Why is the concentration of water considered constant in the K_w expression?

Table: K_w at various temperature.

Temp. ($^\circ\text{C}$)	Value of K_w
0	0.11×10^{-14}
10	0.30×10^{-14}
25	1.0×10^{-14}
40	3.00×10^{-14}
100	7.5×10^{-14}

CHEMISTRY NOTES FOR CLASS - 11 (FOR PUNJAB)

In case of addition of small amount of acid,

$$[H^+] > [OH^-]$$

While in the case of addition of few drops of a base,

$$[OH^-] > [H^+]$$

- During both of these additions, the values of K_w will remain the same, i.e. 10^{-14} at 25°C .

pH AND pOH

In 1909, Sorenson, a Danish biochemist, introduced the term pH and pOH. So, the scales of pH and pOH were developed.

➤ **pH:**

"The negative logarithm of hydrogen ion concentration in the solution is called pH."

$$pH = -\log[H^+]$$

➤ **pOH:**

"The negative logarithm of hydroxide ion concentration in the solution is called pOH."

$$pOH = -\log[OH^-]$$

For neutral water, $pH = -\log[10^{-7}] = 7$

$$pOH = -\log[10^{-7}] = 7$$

Also, $pH + pOH = 14$

The value of pH normally varies between $0 \rightarrow 14$ at 25°C .

The pH values of some familiar aqueous solutions are shown in Figure. This table can help you to understand the acidic or basic nature of commonly used solutions.




Figure: pH Values of some common substances

Sample Problem 9.1

In a solution, the pH is 9.2. Determine the ionic product of water K_w at 25°C .

Solution:

Calculate the pOH from the pH:

$$pOH = 14 - pH$$

$$pOH = 14 - 9.2$$

$$pOH = 4.8$$

Find the concentration of H^+ and OH^- ,

Taking antilog of pH and pOH

$$\text{antilog } pH = \text{antilog}$$

$$[OH^-] = 10^{-pOH}$$

CHEMISTRY NOTES FOR CLASS - 11 (FOR PUNJAB)

$$[H^+] = 10^{-9.2}$$

$$[OH^-] \approx 6.3 \times 10^{-10} M$$

Similarly, antilog pOH = antilog (4.8)

$$[OH^-] = 10^{-pOH}$$

$$[OH^-] = 10^{-4.8}$$

$$[OH^-] \approx 1.6 \times 10^{-5} M$$

Sample Problem 9.2

The ionic product of water at a certain temperature is $K_w = 1.0 \times 10^{-14}$ at $25^\circ C$. If the concentration of H^+ ions in a solution is $1.0 \times 10^{-7} M$.

Calculate the concentration of OH^- ions, the pH and pOH of the solution.

Solution:

Use the expression for K_w :

$$K_w = [H^+][OH^-]$$

Substitute the known values of K_w and the concentration of H^+ :

$$1.0 \times 10^{-14} = (1.0 \times 10^{-7})[OH^-]$$

Solve for the concentration of OH^-

$$[OH^-] = \frac{1.0 \times 10^{-14}}{1.0 \times 10^{-7}} = 1.0 \times 10^{-7}$$

Calculate the pH:

$$\begin{aligned} pH &= -\log[H^+] \\ &= -\log(1.0 \times 10^{-7}) = 7 \end{aligned}$$

Calculate the pOH:

$$\begin{aligned} pOH &= -\log[OH^-] \\ &= -\log(1.0 \times 10^{-7}) \\ &= 7 \end{aligned}$$

QUICK CHECK 9.2


- a) A solution is prepared by mixing equal volumes of two solutions: one with a pH of 4.0 and another with a pH of 10.0. Calculate the K_w for this mixture at $25^\circ C$.

Ans. Given:

- **Solution A:** pH = 4.0 $\rightarrow [H^+] = 10^{-4} M$
- **Solution B:** pH = 10.0 $\rightarrow [OH^-] = 10^{-4} M$ (since pH 10 $\Rightarrow$ pOH 4).

Step 1: Mix Equal Volumes

- $[H^+]$ from A and $[OH^-]$ from B neutralize each other:



- **Result:** $[H^+] = [OH^-] = 10^{-4} M$ cancel out, leaving neutral water (pH 7).

Step 2: Final pH

- The mixture is neutral (pH = 7.0) because the H^+ and OH^- concentrations were equal and fully reacted.

Note on K_w :

- $K_w = [H^+][OH^-] = 1.0 \times 10^{-14}$ (constant at $25^\circ C$) for all aqueous solutions, including this mixture.

Final Answer:

The mixture has:

- pH = 7.0 (neutral).
- $k = 1.0 \times 10^{-14}$ (at $25^\circ C$).

CHEMISTRY NOTES FOR CLASS - 11 (FOR PUNJAB)

- b) At a specific temperature, the ionic product of water K_w is 1.0×10^{-14} . If the concentration of OH^- ions in a solution is $2.5 \times 10^{-8} \text{ M}$. Calculate the concentration of H^+ ions and the pH of the solution.

Ans. Given:

$$K_w = 1.0 \times 10^{-14} \text{ (at the given temperature).}$$

$$[\text{OH}^-] = 2.5 \times 10^{-8} \text{ M}$$

Step 1: Calculate $[\text{H}^+]$

Using the ionic product of water:

$$K_w = [\text{H}^+][\text{OH}^-]$$

$$[\text{H}^+] = \frac{K_w}{[\text{OH}^-]} = \frac{1.0 \times 10^{-14}}{2.5 \times 10^{-8}} = 4.0 \times 10^{-7} \text{ M}$$

Step 2: Calculate pH

$$\text{pH} = -\log [\text{H}^+] = -\log (4.0 \times 10^{-7})$$

$$\text{pH} \approx 6.40 \text{ (since } \log(4.0) \approx 0.60 \text{ and } \log(10^{-7}) = -7 \text{)}$$

Final Answer:

$$[\text{H}^+] = 4.0 \times 10^{-7} \text{ M}$$

$$\text{pH} = 6.40$$

- c) Copper-plate etching solutions is prepared by diluting concentrated HNO_3 to 0.30 M HNO_3 . Calculate $[\text{H}^+]$, pH, $[\text{OH}^-]$ and pOH of this solutions at 25°C .

Ans. Given:

- HNO_3 concentration: 0.30 M (a strong acid, fully dissociates).
- At 25°C : $K_w = 1.0 \times 10^{-14}$.

1. $[\text{H}^+]$ and pH:

Since HNO_3 is a strong acid:

$$[\text{H}^+] = 0.30 \text{ M}$$

$$\text{pH} = -\log [\text{H}^+] = -\log(0.30) \approx 0.52$$

2. $[\text{OH}^-]$ and pOH:

Using $K_w = [\text{H}^+][\text{OH}^-]$:

$$[\text{OH}^-] = \frac{K_w}{[\text{H}^+]} = \frac{1.0 \times 10^{-14}}{0.30} = 3.3 \times 10^{-14} \text{ M}$$

$$\text{pOH} = -\log [\text{OH}^-] = -\log (3.3 \times 10^{-14}) \approx 13.48$$

Final Results:

$$[\text{H}^+] = 0.30 \text{ M}$$

$$\text{pH} = 0.52$$

$$[\text{OH}^-] = 3.3 \times 10^{-14} \text{ M}$$

$$\text{pOH} = 13.48$$

- HNO_3 fully dissociates, so $[\text{H}^+]$ equals its initial concentration.
- $\text{pH} + \text{pOH} = 14$ (verified: $0.52 + 13.48 = 14$).
- Highly acidic ($\text{pH} \ll 7$) due to concentrated HNO_3 .

IONIZATION CONSTANTS OF ACIDS (K_a)

Acids and bases, when dissolved in water, may or may not be completely dissociated. Many acids are weak electrolytes and ionize to an extent which is much less than 100 %. The value of K_a called the dissociation constant of an acid, is the quantitative measure of the strength of the acid.

➤ Mathematical Derivation of K_a

Suppose we have an acid HA dissolved in water,



Rack Your Mind!

5. What is K_a in acid-base chemistry?

CHEMISTRY NOTES FOR CLASS - 11 (FOR PUNJAB)

302

K_c for the reversible reaction will be written as follows:

$$K_c = \frac{[H_3O^+][A^-]}{[HA][H_2O]}$$

$$K_c[H_2O] = \frac{[H_3O^+][A^-]}{[HA]}$$

Let $K_c[H_2O] = K_a$

Hence

$$K_a = \frac{[H_3O^+][A^-]}{[HA]}$$

This equation can be used to calculate K_a for any acidic solution if we know the pH or $[H^+]$ of that solution and the initial concentration of the dissolved acid $[HA]$. This can also be used to calculate the equilibrium concentration of H_3O^+ and A^- produced if we know the initial concentration of acid HA and its K_a value.

➤ K_a and Strength of Acids:

When $K_a < 10^{-3}$ acid is weak.

$K_a = 1$ to 10^{-3} acid is moderately strong.

$K_a > 1$ acid is strong.

The values of K_a for some acids are given in Table.


Table: The values of K_a for some acids

Acid	Dissociation	K_a	Relative strength
HCl	$HCl \rightleftharpoons H^+ + Cl^-$	very large (10^{-2})	Very strong
HNO ₃	$HNO_3 \rightleftharpoons H^+ + NO_3^-$	very large (10^{-5})	Very strong
H ₂ SO ₄	$H_2SO_4 \rightleftharpoons H^+ + HSO_4^-$	Large (10^{-2})	Very strong
HSO ₄ ⁻	$HSO_4^- \rightleftharpoons H^+ + SO_4^{2-}$	1.3×10^{-4}	Strong
HF	$HF \rightleftharpoons H^+ + F^-$	6.7×10^{-5}	Weak
CH ₃ COOH	$CH_3COOH \rightleftharpoons H^+ + CH_3COO^-$	1.85×10^{-5}	Weak
H ₂ CO ₃	$H_2CO_3 \rightleftharpoons H^+ + HCO_3^-$	4.4×10^{-7}	Weak
H ₂ S	$H_2S \rightleftharpoons H^+ + HS^-$	1.0×10^{-7}	Weak
NH ₄ ⁺	$NH_4^+ \rightleftharpoons H^+ + NH_3$	5.7×10^{-10}	Weak
HCO ₃ ⁻	$HCO_3^- \rightleftharpoons H^+ + CO_3^{2-}$	4.7×10^{-12}	Weak
H ₂ O	$H_2O \rightleftharpoons H^+ + OH^-$	1.8×10^{-15}	Very weak


➤ Calculating H_3O^+ ions from K_a

The weak acids do not completely dissociate in water. Therefore, the concentration of hydronium ions is not equal to the initial concentration of the acid. So, the equilibrium expression is used to calculate the H_3O^+ concentration.


For a weak acid HA , the dissociation in water can be represented as:



In water, H^+ combines with water to form H_3O^+ :




So, the equilibrium equation in terms of $H_3O^+_{(aq)}$ is:



CHEMISTRY NOTES FOR CLASS - 11 (FOR PUNJAB)

Let the initial concentration of the weak acid HA be $C \text{ mol dm}^{-3}$.
At equilibrium:



The expression for the acid dissociation constant (K_a) is:

$$K_a = \frac{[\text{H}_3\text{O}^+][\text{A}^-]}{[\text{HA}]}$$

Substitute the equilibrium concentrations into this expression:

$$K_a = \frac{x \cdot x}{C - x} = \frac{x^2}{C - x}$$

For weak acids, x is usually very small compared to C , so $C - x \approx C$. Therefore, the expression simplifies to:

$$K_a \approx \frac{x^2}{C}$$

Taking the square root of both sides:

$$x \approx \sqrt{K_a \times C}$$

Thus, the concentration of H_3O^+ is given by:

$$x \approx \sqrt{K_a \times C}$$

Sample Problem 9.3

Calculate the concentration of $[\text{H}_3\text{O}^+]$ in a $0.1 \text{ M (mol dm}^{-3})$ solution of acetic acid $[\text{CH}_3\text{COOH}]$, given that the acid dissociation constant K_a is 1.8×10^{-5} .

Solution:

$$[\text{H}_3\text{O}^+] \approx \sqrt{K_a \times C}$$

$$[\text{H}_3\text{O}^+] \approx \sqrt{1.8 \times 10^{-5} \times 0.1}$$

$$[\text{H}_3\text{O}^+] \approx \sqrt{0.18 \times 10^{-6}}$$

$$[\text{H}_3\text{O}^+] \approx 1.34 \times 10^{-3} \text{ M (mol dm}^{-3})$$

The concentration $[\text{H}_3\text{O}^+]$ in the solution is approximately $1.34 \times 10^{-3} \text{ mol dm}^{-3}$.



QUICK CHECK 9.3

a) The pH of a 0.10 M solution of formic acid, HCOOH , at 25°C is 2.38 . Calculate K_a for formic acid at this temperature.

Ans. Given:

- Concentration of HCOOH ($[\text{HCOOH}]$): 0.10 M
- pH of the solution: 2.38
- Temperature: 25°C

Step 1: Calculate $[\text{H}^+]$ from pH

$$[\text{H}^+] = 10^{-\text{pH}} = 10^{-2.38} \approx 4.17 \times 10^{-3} \text{ M}$$

Step 2: Set Up the Equilibrium Expression

Formic acid dissociates as:



At equilibrium:

- $[\text{H}^+] = [\text{HCOO}^-] = 4.17 \times 10^{-3} \text{ M}$
- $[\text{HCOOH}] \approx 0.10 - 4.17 \times 10^{-3} \approx 0.0958 \text{ M}$ (since $4.17 \times 10^{-3} \ll 0.10$, the approximation is valid).

Step 3: Calculate K_a

$$K_a = \frac{[\text{H}^+][\text{HCOO}^-]}{[\text{HCOOH}]} = \frac{(4.17 \times 10^{-3})^2}{0.0958} \approx 1.81 \times 10^{-4}$$

Final Answer:

$$K_a \approx 1.81 \times 10^{-4}$$

CHEMISTRY NOTES FOR CLASS - 11 (FOR PUNJAB)

- b) Calculate the concentration of $[H_3O^+]$ in a 0.1 M solution of nitrous acid $[HNO_2]$, given that the acid dissociation constant K_a is 4×10^{-4} .

Ans. Step 1: Write the Dissociation Equation and K_a Expression



$$K_a = \frac{[H^+][NO_2^-]}{[HNO_2]} = 4 \times 10^{-4}$$

Step 2: Define Equilibrium Concentrations

Let $x = [H^+] = [NO_2^-]$ at equilibrium.

Since HNO_2 is a weak acid, its dissociation is small, so:

$$[HNO_2] \approx 0.1 - x \approx 0.1 \text{ M (assuming } x \ll 0.1)$$

Step 3: Substitute into K_a and Solve for x

$$K_a = \frac{x \cdot x}{0.1} = 4 \times 10^{-4}$$

$$x^2 = 4 \times 10^{-5}$$

$$x = \sqrt{4 \times 10^{-5}} = 2 \times 10^{-2.5} \approx 6.32 \times 10^{-3} \text{ M}$$

Step 4: Verify the Approximation

Check if $x \ll 0.1$:

$$\frac{6.32 \times 10^{-3}}{0.1} = 6.32\% (<5\% \text{ is ideal, but } <10\% \text{ is acceptable})$$

The approximation holds reasonably well. For higher precision, solve the quadratic equation:

$$(x^2 + 4 \times 10^{-4})(x - 4 \times 10^{-5}) = 0$$

Using the quadratic formula:

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-4 \times 10^{-5} \pm \sqrt{(4 \times 10^{-5})^2 + 1.6 \times 10^{-4}}}{2}$$

$$x \approx 6.16 \times 10^{-3} \text{ M (minor correction)}$$

Final Answer:

$$[H_3O^+] = 6.32 \times 10^{-3} \text{ M (or } 6.16 \times 10^{-3} \text{ M with quadratic)}$$

- c) A vinegar sample is found to have 0.837 M CH_3COOH . Its hydronium ion concentration is found to be $3.86 \times 10^{-3} \text{ mol dm}^{-3}$. Calculate K_a for acetic acid.

Ans. Given:

- Initial concentration of CH_3COOH ($[CH_3COOH]$): 0.837 M
- Hydronium ion concentration ($[H_3O^+]$ or $[H_3O^+]$): $3.86 \times 10^{-3} \text{ M}$

Step 1: Write the Dissociation Equation

Acetic acid partially dissociates in water:



Step 2: Determine Equilibrium Concentrations

At equilibrium:

- $[H_3O^+] = [CH_3COO^-] = 3.86 \times 10^{-3} \text{ M}$ (from dissociation).
- $[CH_3COOH] \approx 0.837 - 3.86 \times 10^{-3} \approx 0.833 \text{ M}$. (Approximation valid since $3.86 \times 10^{-3} \ll 0.837$)

Step 3: Calculate K_a

Using the equilibrium expression:

$$K_a = \frac{[H_3O^+][CH_3COO^-]}{[CH_3COOH]} = \frac{(3.86 \times 10^{-3})^2}{0.833}$$

$$K_a = \frac{1.49 \times 10^{-5}}{0.833} \approx 1.79 \times 10^{-5}$$

Final Answer:

$$K_a = 1.79 \times 10^{-5}$$

CHEMISTRY NOTES FOR CLASS - 11 (FOR PUNJAB)

COMMON ION EFFECT

[Exercise L.Q.5]

20

Q. Discuss applications and implications of the common ion effect in various fields.


Definition:

"The suppression of ionization of a weak electrolyte by adding a common ion to it is called common ion effect".


➤ Applications of Common Ion Effect

➤ Purification of NaCl:

NaCl is purified by passing hydrogen chloride gas through the saturated brine. Sodium chloride is fully ionized in the solution. Equilibrium constant expression for this process can be written as follows:



HCl also ionizes in solution.




On passing HCl gas, concentration of Cl⁻ ions is increased. Therefore, NaCl crystallizes out of the solution to maintain the constant value of equilibrium constant. This type of effect is called common ion effect.

The addition of a common ion to the solution of less soluble electrolyte suppresses its ionization and the concentration of unionized species increases, which may come out as a precipitate.



➤ Precipitation of KClO₃ by KCl:

The solubility of a less soluble salt KClO₃ in water is suppressed by the addition of more soluble salt KCl by common ion effect. K⁺ is a common ion. The ionization of KClO₃ is suppressed and it settles down as precipitate.



➤ Precipitation of 2nd Group Basic Radicals of Salt Analysis:

The dissociation of a weak acid H₂S in water can be suppressed by the addition of stronger acid HCl. H⁺ is a common ion. H₂S becomes less dissociated in acidic solution. In this way, low concentration of S²⁻ is developed.




This low concentration of S²⁻ ions helps to do the precipitation of radicals of second group basic radicals during salt analysis.



➤ Precipitation of 3rd Group Basic Radicals of Salt Analysis:

An addition of NH₄Cl in NH₄OH solution suppresses the concentration of OH⁻_(aq) due to the presence of a large excess of NH₄⁺ from NH₄Cl. Actually, NH₄Cl is a strong electrolyte. The combination of these two substances is used as reagent in third group basic radicals.



➤ Uses in Qualitative Analysis and Preparation of Buffers:

Common ion effect has its extensive applications in the qualitative analysis and the preparation of buffers.

Note: In common ion effect, the term "Electrolyte" is used instead of acid, base or salt.

Q. Define common ion effect with one example.


- We can suppress the ionization of an electrolyte by common ion effect.

Q. Give applications of common ion effect. (any two)

- Common ion effect is an application of Le-Chatelier's principle.

- In it the term electrolyte is used instead of acid, base or salt.

- In a solution of weak acid, hydrofluoric acid ($K_a = 7.2 \times 10^{-4}$), its salt sodium fluoride produces the common ion (F⁻).



- Common ion effect is always in the reverse direction.



Rack Your Mind!

7. When ionization of a weak electrolyte is suppressed by a strong electrolyte having a common ion is called:
- Hydrolysis
 - Buffering
 - Common ion effect
 - Precipitation reaction

CHEMISTRY NOTES FOR CLASS - 11 (FOR PUNJAB)

BUFFER SOLUTIONS

The solutions which resist the change in their pH when a small amount of an acid or a base is added to them, are called buffer solutions.

- They have a specific constant value of pH. The pH values of buffer solutions do not change on dilution and on keeping for a long time.



Rack Your Mind!

8. What is a buffer solution?


Types of Buffer Solutions

Buffer solutions are mostly prepared by mixing two substances.

- Acidic Buffers:** By mixing a weak acid and its salt with a strong base. Such solutions give acidic buffers with pH less than 7. Mixture of acetic acid and sodium acetate is one of the best examples of acidic buffers.
- Basic Buffers:** By mixing a weak base and its salt with a strong acid. Such solutions will give basic buffers with pH more than 7. Mixture of NH_4OH and NH_4Cl is one of the best examples of basic buffers.

Example:

An acidic buffer consisting of CH_3COOH and CH_3COONa . CH_3COOH being a weak electrolyte undergoes very little dissociation. When CH_3COONa , which is a strong electrolyte, is added to CH_3COOH solution, then the dissociation of CH_3COOH is suppressed, due to common ion effect of CH_3COO^- .



If one goes on adding CH_3COONa in CH_3COOH solution, then the added concentrations of CH_3COO^- decrease the dissociation of CH_3COOH and the pH of solution increases. The Table tells us how the pH value of a mixture of two compounds is maintained. Greater the concentration of acetic acid as compared to CH_3COONa , lesser is the pH of solution.

Table: Effect of addition of acetate ions on the pH of acetic acid solution

$[\text{CH}_3\text{COOH}](\text{mol dm}^{-3})$	$[\text{CH}_3\text{COO}^-](\text{mol dm}^{-3})$	% Dissociation	pH
0.10	0.00	1.3	2.89
0.10	0.05	0.036	4.44
0.10	0.10	0.018	4.74
0.10	0.15	0.012	4.92

Actually, a buffer mentioned above is a large reservoir of CH_3COOH and CH_3COO^- components. When an acid or H_3O^+ ions are added to this buffer, they will react with CH_3COO^- to give back acetic acid and hence the pH of the solution will almost remain unchanged. The reason is that CH_3COOH being a weak acid will prefer to remain undissociated. Similarly, the buffer solution consisting of NH_4Cl and NH_4OH , can resist the change of pH and pOH, when acid or base is added from outside. When a base or OH^- ions are added in it, they will react with H_3O^+ to give back H_2O and the pH of the solution again will remain almost unchanged.

Acidic buffers	Basic buffers
- Those buffers which are formed by the mixing of a weak acid and a salt of it with a strong base are called acidic buffers. - Their pH is usually less than 7. - Henderson's equation for acidic buffers $\text{pH} = \text{pK}_a + \log \frac{[\text{salt}]}{[\text{acid}]}$	- Those buffers which are formed by mixing of a weak base and a salt of it with a strong acid are called basic buffers. - Their pH is more than 7. - Henderson's equation for basic buffers $\text{pOH} = \text{pK}_b + \log \frac{[\text{salt}]}{[\text{base}]}$
e.g. CH_3COOH + CH_3COONa weak acid salt of weak acid	e.g. NH_4OH + NH_4Cl weak base salt of weak base



CHEMISTRY NOTES FOR CLASS - 11 (FOR PUNJAB)

➡ Calculating the pH of a Buffer

Let us discuss preparation of a buffer of definite pH.

Example:

Consider a weak acid HA and its salt NaA with a strong base say NaOH. The reversible reactions for dissociation of HA and NaA are as follows:



Calculation of pH of a buffer solution by using Henderson equation:

$$pH = pK_a - \log \frac{[Acid]}{[Salt]}$$

Interchanging the numerator and denominator, the sign of log changes.

$$pH = pK_a + \log \frac{[Salt]}{[Acid]}$$

This relationship is called **Henderson's equation**.

Factors Affecting pH of a Buffer:

Henderson's equation shows that two factors which govern the pH of a buffer solution are as follows:

- (i) pK_a of the acid used.
- (ii) The ratio of the concentrations of the salt and the acid.
 - The best buffer is prepared by taking equal concentration of salt and acid.
 - So, pH is controlled by pK_a of the acid.

Example: For acetic acid - sodium acetate buffer, if

$$\begin{aligned} [CH_3COOH] &= [CH_3COONa] \\ pK_a &= 4.74 \end{aligned}$$

Then
$$pH = pK_a + \log \frac{[CH_3COONa]}{[CH_3COOH]}$$

$$pH = pK_a + \log(1)$$

So,
$$pH = pK_a + 0 = pK_a$$

$$pH = 4.74$$

It means that the pH of this buffer is just equal to the pK_a of the acid.

Interesting Information!

The body's blood buffering system, involving bicarbonates, helps maintain a



QUICK CHECK 9.4

a) Explain the impact of common ion effect on solubility.

Ans. Effect of Common Ion on Solubility

The presence of a common ion decreases the solubility of a slightly soluble ionic compound. Consider a saturated solution of $PbCrO_4$, which is a sparingly soluble ionic salt.



Now add Na_2CrO_4 which is a soluble salt. CrO_4^{2-} is the common ion. It combines with Pb^{2+} to form more insoluble $PbCrO_4$. So equilibrium is shifted to the left to keep K_{sp} constant.

b) To a saturated solution of $AgCl$, some of $NaCl$ solution was added.

i. State the effect of the concentration of Ag^+ on the equilibrium.

Ans. Effect on $[Ag^+]$:

The addition of $NaCl$ (which provides Cl^- ions) to a saturated $AgCl$ solution decreases the concentration of Ag^+ ions at equilibrium.

CHEMISTRY NOTES FOR CLASS - 11 (FOR PUNJAB)

ii. Explain your answer with respect to the common ion effect.

Ans. Explanation (Common Ion Effect):

Equilibrium Shift:

- The dissolution equilibrium of AgCl is:



- Adding NaCl introduces excess Cl^- , shifting the equilibrium left (Le-Chatelier's Principle) to reduce the added Cl^- , causing more AgCl to precipitate.

Mathematical Impact:

- The solubility product (K_{sp}) must remain constant:

$$K_{sp} = [\text{Ag}^+][\text{Cl}^-]$$

If $[\text{Cl}^-]$ increases, $[\text{Ag}^+]$ must decrease to maintain K_{sp} .

Result:

- Solubility of AgCl decreases, and the solution becomes less saturated with Ag^+ ions.

c) How does a buffer maintain pH stability?

Ans. A buffer maintains pH stability by resisting changes in pH when small amounts of acid (H^+) or base (OH^-) are added. Here's how it works concisely:

Mechanisms:

1. Composition:

- A buffer consists of a weak acid (HA) and its conjugate base (A^-) (e.g., $\text{CH}_3\text{COOH}/\text{CH}_3\text{COO}^-$) or a weak base (B) and its conjugate acid (BH^+) (e.g., $\text{NH}_3/\text{NH}_4^+$).

2. Action Against Added Acid (H^+):

- The conjugate base (A^-) neutralizes H^+ :



- Example: In $\text{CH}_3\text{COO}^-/\text{CH}_3\text{COOH}$ buffer, CH_3COO^- consumes H^+ to form CH_3COOH .

3. Action Against Added Base (OH^-):

- The weak acid (HA) neutralizes OH^- :



- Example: CH_3COOH donates H^+ to OH^- , forming water and CH_3COO^- .

4. pH Stability:


- The equilibrium between HA and A^- ensures $[\text{H}^+]$ (and thus pH) changes minimally.
- The Henderson-Hasselbalch equation predicts buffer pH:

$$\text{pH} = \text{pK}_a + \log\left(\frac{[\text{A}^-]}{[\text{HA}]}\right)$$

Example:

A phosphate buffer ($\text{H}_2\text{PO}_4^-/\text{HPO}_4^{2-}$) maintains blood pH (~7.4) by:

- Neutralizing added H^+ :



- Neutralizing added OH^- :



Conclusion:

Buffers stabilize pH by shifting equilibrium to consume added H^+ or OH^- , keeping the $[\text{HA}]/[\text{A}^-]$ ratio (and thus pH) nearly constant.

d) Calculate the pH of a buffer consisting of 0.50 M HF and 0.45 M of a fluoride (F^-) salt before and after addition of 0.40 g NaOH to 1.0 dm³ of the buffer (K_a of HF = 6.8×10^{-4}).

Ans. Given Buffer Solution:

$$[\text{HF}] = 0.50 \text{ M}$$

$$[\text{F}^-] = 0.45 \text{ M}$$

$$\text{Volume} = 1.0 \text{ dm}^3$$

$$K_a \text{ of HF} = 6.8 \times 10^{-4}$$

- Added NaOH = 0.40 g (Molar mass = 40 g/mol $\rightarrow$ 0.01 mol NaOH in 1.0 dm³ $\rightarrow$ $[\text{OH}^-] = 0.01 \text{ M}$)

CHEMISTRY NOTES FOR CLASS - 11 (FOR PUNJAB)

Step 1: Calculate Initial pH (Before NaOH Addition)

Using the Henderson-Hasselbalch equation:

$$\text{pH} = \text{pK}_a + \log\left(\frac{[\text{F}^-]}{[\text{HF}]}\right)$$


$$\text{pK}_a = -\log(6.8 \times 10^{-4}) \approx 3.17$$

$$\text{pK}_a = -\log(6.8 \times 10^{-4}) \approx 3.17$$

$$\text{pH} = 3.17 + \log(0.450/0.50) = 3.17 + \log(0.9) \approx 3.17 - 0.046 = 3.12$$

Step 2: Calculate pH After Adding NaOH (0.01 M OH⁻)

NaOH reacts with HF (weak acid) in the buffer:



• HF consumed: $0.50 - 0.01 = 0.49 \text{ M}$

• F⁻ produced: $0.45 + 0.01 = 0.46 \text{ M}$

New pH:

$$\text{pH} = 3.17 + \log(0.46/0.49) = 3.17 + \log(0.94) \approx 3.17 - 0.027 = 3.14$$

e) Calculate the pH of a buffer solution in which 0.11 molar CH₃COONa and 0.09 molar acetic acid solutions are present. K_a for CH₃COOH is 1.85×10^{-5} .

Given Data:

$$[\text{CH}_3\text{COONa}] = 0.11 \text{ M}$$

$$[\text{CH}_3\text{COOH}] = 0.09 \text{ M}$$

$$K_a \text{ of } \text{CH}_3\text{COOH} = 1.85 \times 10^{-5}$$

Required:

$$\text{pH of Buffer solution} = ?$$

Solution:

$$\text{pK}_a = -\log K_a$$

$$= -\log 1.8 \times 10^{-5} = 4.74$$

$$\text{pH} = \text{pK}_a + \log \frac{[\text{Salt}]}{[\text{Acid}]}$$

$$\text{pH} = 4.74 + \log \frac{0.11}{0.09}$$

$$\text{pH} = 4.74 + 0.087$$

$$\text{pH} = 4.83$$

SOLUBILITY PRODUCT

(For slightly soluble ionic compounds)

Q. What is the solubility product for sparingly soluble salts? Give its two applications.

(Exercise L.Q.6)

Definition:

"The solubility product is the product of the concentrations of ions raised to an exponent equal to the coefficient of the balanced equation".


OR

"The product of the equilibrium concentration of ions raised to an exponent equal to the coefficient of the balanced equation at given temperature is called solubility product (K_{sp})".

When a soluble ionic compound is dissolved in water, like NaCl, it dissociates completely into ions. But for slightly soluble salts the dissociation is not complete at equilibrium stage.

➤ **Example 1, Lead Sulphate:**

When Lead Sulphate (PbSO₄) is shaken with water the solution contains Pb²⁺, SO₄²⁻ and undissociated PbSO₄. It means that equilibrium exists between solid solute, PbSO₄ and the dissolved ions, SO₄²⁻ and Pb²⁺. Lead sulphate is a well-known sparingly soluble compound and it dissociates to a very small extent like PbCl₂.



$$K_c = \frac{[\text{Pb}^{2+}][\text{SO}_4^{2-}]}{[\text{PbSO}_4]}$$

CHEMISTRY NOTES FOR CLASS - 11 (FOR PUNJAB)

310

Being a sparingly soluble salt, the concentration of lead sulphate (PbSO_4) almost remains constant. Bring $[\text{PbSO}_4]$ on L.H.S. with K_c .

$$\begin{aligned} K_c [\text{PbSO}_4] &= [\text{Pb}^{2+}][\text{SO}_4^{2-}] \\ \text{if } K_c [\text{PbSO}_4] &= K_{sp} \\ \text{then } K_{sp} &= [\text{Pb}^{2+}][\text{SO}_4^{2-}] = 1.6 \times 10^{-8} \text{ at } 25^\circ \text{C} \end{aligned}$$

K_{sp} is called the solubility product of PbSO_4 . It is the product of molar solubilities of two ions at equilibrium stage.

➤ Example 2, Lead Chloride:

Consider a sparingly soluble salt, PbCl_2 . $K_{sp} = [\text{Pb}^{2+}][\text{Cl}^-]^2$

K_{sp} is usually a very small quantity at room temperature. The value of K_{sp} is temperature dependent. The value of K_{sp} is a measure of the dissociation of sparingly soluble salt.

The following Table shows the K_{sp} values of some slightly soluble ionic compounds.

Table: K_{sp} values for some ionic compounds

Salt	Ion Product	K_{sp}	Salt	Ion Product	K_{sp}
AgBr	$[\text{Ag}^+][\text{Br}^-]$	5.0×10^{-13}	CuS	$[\text{Cu}^{2+}][\text{S}^{2-}]$	8×10^{-34}
AgCl	$[\text{Ag}^+][\text{Cl}^-]$	1.8×10^{-10}	Fe_2S_3	$[\text{Fe}^{3+}]^2[\text{S}^{2-}]^3$	1.4×10^{-85}
$\text{Al}(\text{OH})_3$	$[\text{Al}^{3+}][\text{OH}^-]^3$	3×10^{-34}	MgCO_3	$[\text{Mg}^{2+}][\text{CO}_3^{2-}]$	3.5×10^{-8}
BaSO_4	$[\text{Ba}^{2+}][\text{SO}_4^{2-}]$	1.1×10^{-10}	MnS	$[\text{Mn}^{2+}][\text{S}^{2-}]$	3×10^{-11}
CaCO_3	$[\text{Ca}^{2+}][\text{CO}_3^{2-}]$	3.3×10^{-9}	PbCrO_4	$[\text{Pb}^{2+}][\text{CrO}_4^{2-}]$	2.3×10^{-13}
CaF_2	$[\text{Ca}^{2+}][\text{F}^-]^2$	3.2×10^{-11}	PbSO_4	$[\text{Pb}^{2+}][\text{SO}_4^{2-}]$	1.6×10^{-8}

➤ Applications of Solubility Product

a) Determination of Solubility from K_{sp}

For this purpose, we need the formula of the compound and K_{sp} value. Then the unknown molar solubility S is calculated and the concentration of the ions is determined. Table shows the relationship between the K_{sp} values and the solubility of some sparingly soluble compounds.

Table: The relation between solubility and solubility products of some salts

Formula	No. of Ions	K_{sp}	Solubility g dm^{-3}
PbSO_4	2	1.69×10^{-4}	1.3×10^{-4}
$\text{Ca}(\text{OH})_2$	3	6.5×10^{-6}	1.175×10^{-2}
CaF_2	3	3.2×10^{-11}	2.0×10^{-4}
Ag_2CrO_4	3	2.6×10^{-12}	8.7×10^{-5}

Rack Your Mind!

9. What is the effect of temperature on the K_{sp} value?

- A) It always remains constant
- B) It decreases with temperature
- C) It changes with temperature
- D) It increases with pressure

CHEMISTRY NOTES FOR CLASS - 11 (FOR PUNJAB)

Smart  Thinking				
Relationship Between K_{sp} and Molar Solubility (s)				
Compound	Cation	Anion	Ksp Expression	Relation Between Ksp and s
AgCl	s	s	$[Ag^+][Cl^-]$	$K_{sp} = s^2 \Rightarrow s = (K_{sp})^{\frac{1}{2}}$
BaSO ₄	s	s	$[Ba^{2+}][SO_4^{2-}]$	$K_{sp} = s^2 \Rightarrow s = (K_{sp})^{\frac{1}{2}}$
PbF ₂	s	2s	$[Pb^{2+}][F^-]^2$	$K_{sp} = 4s^3 \Rightarrow s = \left(\frac{K_{sp}}{4}\right)^{\frac{1}{3}}$
Ag ₂ CO ₃	2s	s	$[Ag^+]^2[CO_3^{2-}]$	$K_{sp} = 4s^3 \Rightarrow s = \left(\frac{K_{sp}}{4}\right)^{\frac{1}{3}}$
Al(OH) ₃	s	3s	$[Al^{3+}][OH^-]^3$	$K_{sp} = 27s^4 \Rightarrow s = \left(\frac{K_{sp}}{27}\right)^{\frac{1}{4}}$
Ca ₃ (PO ₄) ₂	3s	2s	$[Ca^{2+}]^3[PO_4^{3-}]^2$	$K_{sp} = 108s^5 \Rightarrow s = \left(\frac{K_{sp}}{108}\right)^{\frac{1}{5}}$



Sample Problem 9.4

Ca(OH)₂ is a sparingly soluble compound. Its solubility product is 6.5×10^{-6} . Calculate the solubility of Ca(OH)₂.

Solution:

Let the solubility be represented by S in terms of mol dm^{-3} .

The balanced equation is:



The concentration of OH⁻ is double than the concentration of Ca²⁺, so

$$K_{sp} = [Ca^{2+}][OH^-]^2 = S \times (2S)^2$$

$$4S^3 = 6.5 \times 10^{-6}$$

$$S^3 = \frac{6.5 \times 10^{-6}}{4}$$

$$S = (1.625)^{1/3} \times 10^{-2}$$

$$S = 1.18 \times 10^{-2} \text{ mol dm}^{-3}$$


Hence, at equilibrium stage $1.18 \times 10^{-2} \text{ mol dm}^{-3}$ of Ca²⁺ and $2 \times 1.18 \times 10^{-2} \text{ mol dm}^{-3} = 2.36 \times 10^{-2} \text{ mol dm}^{-3}$ OH⁻ are present in the solution. In this way, we have calculated the individual concentrations of Ca²⁺ and OH⁻ ions from the solubility product of Ca(OH)₂.

b) Common Ion Effect

The presence of a common ion decreases the solubility of a slightly soluble ionic compound. In order to explain it, consider a saturated solution of PbCrO₄, which is a sparingly soluble ionic salt.

$\text{Solubility} \propto \frac{1}{\text{concentration (no.) of common ions}}$

CHEMISTRY NOTES FOR CLASS - 11 (FOR PUNJAB)



Now add Na_2CrO_4 which is a soluble salt.



CrO_4^{2-} is the common ion. It combines with Pb^{2+} to form more insoluble PbCrO_4 . So equilibrium is shifted to the left to keep K_{sp} constant.

c) Predicting Precipitation

The solubility product (K_{sp}) helps in predicting whether the precipitation of a salt will occur or not. For example, the solubility product of CaSO_4 is 2×10^{-5} . If we add $10^{-2} \text{ mol dm}^{-3}$ solution Ca^{2+} to $10^{-2} \text{ mol dm}^{-3}$ solution of SO_4^{2-} ions at 25°C . The concentrations of each ionic species can be calculated as:

Ionic Product	Type of Solution	Precipitation
$> K_{sp}$	Supersaturated	Yes
$= K_{sp}$	Saturated	No
$< K_{sp}$	Unsaturated	No

$$[\text{Ca}^{2+}] = [\text{SO}_4^{2-}] = \frac{10^{-2}}{2} = 5.0 \times 10^{-3} \text{ mol dm}^{-3}$$

$$\begin{aligned} [\text{Ca}^{2+}][\text{SO}_4^{2-}] &= 5.0 \times 10^{-3} \text{ mol dm}^{-3} \times 5.0 \times 10^{-3} \text{ mol dm}^{-3} \\ &= 2.5 \times 10^{-5} \text{ mol}^2 \text{ dm}^{-6} \\ &= 2.5 \times 10^{-5} \text{ mol}^2 \text{ dm}^{-6} > K_{sp} \text{ of } \text{CaSO}_4 \end{aligned}$$

As the product of concentrations is greater than K_{sp} , therefore CaSO_4 will precipitate out.

☑ QUICK CHECK 9.5

- a) The solubility product constant (K_{sp}) of silver chloride (AgCl) is 1.77×10^{-10} at 25°C . A solution already contains 0.10 M of sodium chloride (NaCl). Find the solubility of AgCl in a solution that contains 0.10 NaCl .

Ans. Given:

- K_{sp} of $\text{AgCl} = 1.77 \times 10^{-10}$
- $[\text{Cl}^-]$ from $\text{NaCl} = 0.10 \text{ M}$ (this is the common ion).

Step 1: Define Solubility (S) of AgCl

Let S = molar solubility of AgCl (mol/dm^3) in the presence of NaCl .

- Dissolution equation:



- At equilibrium:

$$[\text{Ag}^+] = S (\text{from } \text{AgCl})$$

$$[\text{Cl}^-] = 0.10 \text{ M (from NaCl)} + S = 0.10 \text{ M (since } S \ll 0.10)$$

Step 2: Apply K_{sp} Expression

$$K_{sp} = [\text{Ag}^+][\text{Cl}^-]$$

$$1.77 \times 10^{-10} = S \times 0.10$$

$$S = 1.77 \times 10^{-10} / 0.10 = 1.77 \times 10^{-9} \text{ M}$$

1. Common Ion Effect:

The pre-existing 0.10 M Cl^- from NaCl drastically reduces AgCl solubility (from $1.33 \times 10^{-5} \text{ M}$ in pure water to $1.77 \times 10^{-9} \text{ M}$).

2. Assumption Valid:

$S \ll 0.10$, so $[\text{Cl}^-] \approx 0.10 \text{ M}$ is justified.

Conclusion:

The solubility of AgCl in 0.10 M NaCl is $1.77 \times 10^{-9} \text{ M}$.

CHEMISTRY NOTES FOR CLASS - 11 (FOR PUNJAB)

b) Predict whether CaSO_4 will be precipitated or not when $10^{-3} \text{ mol dm}^{-3}$ of each of Ca^{2+} and SO_4^{2-} is mixed together.

Ans. Given:

$$[\text{Ca}^{2+}] = 10^{-3} \text{ mol dm}^{-3}$$

$$[\text{SO}_4^{2-}] = 10^{-3} \text{ mol dm}^{-3}$$

$$K_{sp} \text{ of } \text{CaSO}_4 = 2.4 \times 10^{-5} \text{ (at } 25^\circ\text{C, literature value).}$$

Step 1: Calculate the Ion Product (Q)

$$Q = [\text{Ca}^{2+}][\text{SO}_4^{2-}] = (10^{-3})(10^{-3}) = 10^{-6}$$

Step 2: Compare Q to K_{sp}

- If $Q > K_{sp}$: Precipitation occurs.
- If $Q < K_{sp}$: No precipitation.

Here: $Q = 10^{-6}$ vs. $K_{sp} = 2.4 \times 10^{-5}$

Since $10^{-6} < 2.4 \times 10^{-5} = 10^{-6} < 2.4 \times 10^{-5}$, $Q < K_{sp}$.

Conclusion:

CaSO_4 will NOT precipitate under these conditions because the ion product (Q) is less than K_{sp} .

c) Solution of potassium carbonate is basic.

i. Explain why the solution is alkaline.

Ans. Solution is Alkaline:


Potassium carbonate (K_2CO_3) dissolves in water to produce CO_3^{2-} , which hydrolyzes to form OH^- ions, making the solution basic:



- CO_3^{2-} (a strong base) reacts with water, accepting protons (H^+) and releasing OH^- .
- K^+ (from K_2CO_3) is a spectator ion and does not hydrolyze.

ii. Also give the equation for the hydrolysis of the above salt.

Ans. Hydrolysis Equation:



The OH^- generated during hydrolysis raises the pH (>7), making the solution alkaline.

Example pH: ~ 11 for 0.1 M K_2CO_3 .



Interesting Information!

The shell of this nautilus is composed mainly of calcium carbonate. The nautilus adjusts conditions so shell material is formed when the concentration of calcium ions and carbonate ions in seawater are high enough to precipitate calcium carbonate.



SALT HYDROLYSIS

When a salt dissolves in water, it dissociates into its constituent ions. These ions can interact with water, affecting the solution's pH depending on the nature of the acid and base from which the salt is derived.


➤ **Hydrolysis:**

"The interaction between salts and water are called hydrolytic reactions and this phenomenon is called hydrolysis".

➤ **Salts of Strong Acids and Strong Bases**

For salts derived from strong acids and strong bases (e.g., sodium chloride, NaCl), the conjugate base of a strong acid (e.g., Cl^- from HCl) is very weak and does not significantly react with water. The conjugate acid of a strong base (e.g., Na^+ from NaOH) is also very weak and does not significantly react with water.

CHEMISTRY NOTES FOR CLASS - 11 (FOR PUNJAB)



Na^+ is the conjugate acid of NaOH (a strong base) and does not affect the pH.

Cl^- is the conjugate base of HCl (a strong acid) and does not affect the pH.

The solution remains neutral.

Examples are KCl ($\text{HCl} + \text{KOH}$), KNO_3 ($\text{HNO}_3 + \text{KOH}$), Na_2SO_4 ($\text{H}_2\text{SO}_4 + \text{NaOH}$) etc.

⇒ Salts of Strong Acids and Weak Bases

For salts derived from strong acids and weak bases (e.g., ammonium chloride, NH_4Cl), The conjugate base of the strong acid (e.g., Cl^-) does not react with water. The conjugate acid of the weak base (e.g., NH_4^+ from NH_3) reacts with water to produce H_3O^+ ions, making the solution acidic.



NH_4^+ is the conjugate acid of NH_3 (a weak base) and reacts with water:



Cl^- is the conjugate base of HCl (a strong acid) and does not affect the pH.

The solution is acidic due to the NH_4^+ ion.

Examples: AlCl_3 , CuSO_4 , NH_4NO_3 etc.

⇒ Salts of Weak Acids and Strong Bases

The example of salt of weak acid and strong base is sodium acetate.



CH_3COO^- is the conjugate base of CH_3COOH (a weak acid) and reacts with water:




Na^+ is the conjugate acid of NaOH (a strong base) and does not affect the pH. The solution is basic due to the CH_3COO^- ion.

Examples: Na_2CO_3 ($\text{H}_2\text{CO}_3 + \text{NaOH}$), Na_3PO_4 ($\text{H}_3\text{PO}_4 + \text{NaOH}$)

⇒ Salts of Weak Acids and Weak Bases

For salts derived from weak acids and weak bases (e.g., ammonium acetate, $\text{NH}_4\text{CH}_3\text{COO}$): The conjugate acid (NH_4^+) and the conjugate base (CH_3COO^-) both affect the pH.



NH_4^+ hydrolyzes to produce H_3O^+ .



CH_3COO^- hydrolyzes to produce OH^- .



The resultant pH of the solution depends on the relative strengths of the conjugate acid and conjugate base. The solution may be acidic, basic, or nearly neutral, depending on which reaction is more dominant.

Examples: NH_4CN ($\text{NH}_4\text{OH} + \text{HCN}$), $(\text{NH}_4)_2\text{CO}_3$ ($\text{H}_2\text{CO}_3 + \text{NH}_3$)

CHEMISTRY NOTES FOR CLASS - 11 (FOR PUNJAB)

ACID-BASE INDICATORS

An indicator is a substance that changes color to mark a titration's endpoint. Acid-base indicators exhibit one color in acid and another in base. Most indicators used in acid-base titration are weak organic acids or weak organic bases. In solution, a weak-acid indicator (HIn) can be represented by the equation below.



In^- is the symbol of the anion part of the indicator. Because the reaction is reversible, both HIn and In^- are present. The colors displayed result from the fact that HIn and In^- have different colors.

In acidic solutions, any In^- ions that are present act as Brønsted bases and accept protons from the acid. The indicator is then present in largely unionized form, HIn. In basic solutions, the OH^- ions from the base combine with the H^+ ions produced by the indicator. The indicator molecules further ionize to equalize the loss of H^+ ions. The indicator is thus present largely in the form of its anion, In^- . The solution now shows the base-indicating color, which for litmus is blue.




Figure: Unionized acid-base indicators and their ionised forms exist in an equilibrium

Different indicators change color at different pH values. The color depends on the relative amounts of HIn and In^- at a given pH. For example, methyl red changes from red to yellow between pH 4.4 and 6.2. At pH 4.4, the indicator exists mostly as HIn molecules, which appear red in the solution. The indicator ranges are given below for some of the indicators commonly used.



Figure: Range and color changes of some common Acid-Base indicators

☞ Selecting a Suitable Indicator

The two general criteria for an indicator to be used in a titration are:

- The pH at the end of the titration should be close to the indicator's neutral point.
- The indicator should indicate a sharp color change near the equivalence point of the titration.

Each pH indicator changes color over a defined range of pH, known as the indicator range. An indicator changes color over a range of about 2 pH units.

(Exercise L.Q.7)

Q. Describe the general shape of a titration curve for a strong acid titrated with a strong base. How can you identify the equivalence point on this titration curve?

☞ Titration Curve and Equivalence Point

A pH curve is a graph of the pH of the solution versus the volume of titrant added. The equivalence point is the point at which the amount of titrant added is just enough to neutralize the analyte solution completely.

Titration curves show how the pH of an acidic or basic solution changes as a basic or acidic solution is added to it. We can use the titration curve to choose an indicator that will show when the titration is complete and we reach the equivalence point. The end point of the titration occurs when the indicator changes color. We choose an indicator with an end point close to the equivalence point.

CHEMISTRY NOTES FOR CLASS - 11 (FOR PUNJAB)

1. Strong Acid-Strong Base Titration Curve:

Consider the titration of HCl with NaOH . Initially, the solution contains only the strong acid. Since the acid is strong, it completely dissociates, leading to a high concentration of H^+ ions and the pH is very low.

As NaOH is added, OH^- ions from NaOH begin to neutralize the H^+ ions from HCl and pH of the solution rises.

The equivalence point is reached when the amount of OH^- added is stoichiometrically equal to the amount of H^+ originally present in the acid. **The pH at the equivalence point in a strong acid-strong base titration is 7.0.**

The titration curve at this stage shows a steep rise in pH, changing quickly from acidic to neutral. The pH value at the endpoint changes about from 4.0 to 10.0. These titrations have a pH of 7.0 at equivalence, so, such phenolphthalein as indicator can be used as they show different colours in this range.




Figure: Strong acid-strong base titration curve

2. Strong Acid-Weak Base Titration Curve

Consider the titration of aqueous NH_3 with HCl .

Initially, the solution contains only the weak base and it only partially dissociates, with a lower concentration of OH^- ions compared to a strong base. The initial pH will be greater than 7 but lower than the pH of a strong base. As HCl is added, the concentration of OH^- decreases, causing a further decrease in pH.

Before reaching the equivalence point, the solution is in a buffer-like region where the pH changes more gradually. The presence of the weak base and its conjugate acid (from the salt formed) creates a buffering effect, which helps to moderate the pH change as the base is added.

The pH at the equivalence point will be less than 7 (pH=5.27) because the conjugate acid slightly dissociates, releasing H^+ ions making the solution acidic ($\text{NH}_4^+ \rightarrow \text{NH}_3 + \text{H}^+$).

After the equivalence point, the pH of the solution decreases rapidly. This is because the strong acid dissociates completely in water, providing a high concentration of hydrogen ions H^+ , which significantly lowers the pH. Methyl orange has its colour change in this range, therefore, it can be used as an indicator for strong acid-weak base titrations.




Figure: strong acid-weak base titration curve



Rack Your Mind!

10. What is the equivalence point in a titration?

Table: pH ranges of common indicators

Indicators	Acid color	Base color	pH Range	Type of Titration
Methyl orange	orange	yellow	3.2 – 4.5	strong acid-strong base strong acid-weak base
Phenolphthalein	colorless	red	8.2 – 10.0	weak acid-strong base

CHEMISTRY NOTES FOR CLASS - 11 (FOR PUNJAB)

QUICK CHECK 9.6

a) Differentiate end point and equivalent point.

Ans. The difference between Endpoint and Equivalence:

Feature	Equivalence Point	Endpoint
Definition	The point where stoichiometrically equivalent amounts of analyte and titrant react.	The point where the indicator changes color, signaling completion of titration.
Theoretical vs. Experimental	Theoretical (calculated from stoichiometry).	Experimental (observed visually or instrumentally).
Dependence on Indicator	Independent of indicator.	Depends on the choice of indicator.
pH (Acid-Base Titration)	pH is determined by the salt formed (e.g., pH = 7 for strong acid-strong base).	pH may slightly differ due to indicator range (e.g., phenolphthalein changes at pH 8.2–10).
Precision	Exact (based on reaction stoichiometry).	Approximate (may not perfectly match equivalence point).

b) Explain how an indicator changes its colour in acidic and basic solution.

Ans. An Indicator Changes Color in Acidic vs. Basic Solutions:

1. Chemical Structure:

- Indicators are **weak acids (HIn)** or **weak bases (InOH)** with different colors in protonated (acidic) and deprotonated (basic) forms.

2. Acidic Solution ($\text{pH} < \text{pK}_a$):

- Excess H^+ ions shift equilibrium left:



- The **protonated form (HIn)** dominates (e.g., red for methyl orange).

3. Basic Solution ($\text{pH} > \text{pK}_a$):

- OH^- ions remove H^+ , shifting equilibrium right:



- The **deprotonated form (In⁻)** dominates (e.g., yellow for methyl orange).

4. Transition Range ($\text{pH} \approx \text{pK}_a \pm 1$):


- Color change occurs (e.g., phenolphthalein turns pink at pH 8.2–10).

Example:

- Methyl Orange:**



- Phenolphthalein:**



Key Idea: The indicator's color reflects the dominant form (HIn or In⁻) based on pH.

c) Suggest a suitable indicator for weak acid and strong base titration.

Ans. For the titration of a weak acid (e.g., CH_3COOH) with a strong base (e.g., NaOH), the best indicator is:

Phenolphthalein

- pH Range:** 8.2–10.0 (colorless → pink).
- Reason:** The equivalence point pH is > 7 (due to hydrolysis of the conjugate base, e.g., CH_3COO^-). Phenolphthalein's range matches this alkaline pH.

CHEMISTRY NOTES FOR CLASS - 11 (FOR PUNJAB)

d) Suggest a suitable indicator for weak acid and weak base titration.

Ans. For weak acid-weak base titrations, no indicator is ideal due to the gradual pH change near the equivalence point. However, if required:

Suggested Indicator: Bromothymol Blue

- **pH Range:** 6.0 (yellow) to 7.6 (blue).
- **Reason:** Covers the neutral pH (≈ 7) typical for weak acid-weak base titrations (e.g., acetic acid + ammonia).

Limitation:

Weak acid-weak base titrations often lack a sharp pH jump, making pH meters more reliable than visual indicators.

Solution File



Rack Your Brain!

Sr. #	Option	Explanation
1.	B	$\text{HCl (acid)} \rightarrow \text{H}^+ + \text{Cl}^-$ (conjugate base) HCl donates a proton, making Cl^- its conjugate base.
2.	D	When an acid donates a proton, it forms its conjugate base, and when a base accepts a proton, it forms its conjugate acid.
3.	S.Q	BF_3 is a Lewis acid because the boron atom has only six electrons in its valence shell, which is less than an octet. This incomplete octet makes BF_3 electron-deficient and able to accept a lone pair from a Lewis base like fluoride ion.
4.	S.Q	Water concentration remains effectively constant (about 55.5 mol dm^{-3}) because water is present in large excess and only a tiny fraction ionizes, so its concentration does not change significantly.
5.	A	$\text{H}^+ = 10^{-\text{pH}}$ $= 10^{-3}$ $= 0.001 \text{ M}$ or $1 \times 10^{-3} \text{ M}$ pH is defined as the negative logarithm of the hydrogen ion concentration.
6.	S.Q	K_a is the acid dissociation constant. It represents the extent to which an acid donates protons in water. A higher K_a value means a stronger acid.
7.	C	Presence of a common ion from a strong electrolyte reduces the ionization of a weak electrolyte, shifting the equilibrium towards the reactants.
8.	S.Q	• A buffer solution resists a change in pH when a small amount of acid or base is added. • It maintains a constant pH and is stable over time and on dilution. • Buffers are made from a weak acid/base and its salt.
9.	C	Based on "Le Chatelier's Principle" the system shifts to oppose the temperature change. So "temperature can increase or decrease K_{sp} " depending on the heat change of the dissolution reaction.
10.	S.Q	• The equivalence point is when the amount of acid equals the amount of base. • It is where complete neutralization occurs. • It is not always at pH 7, depending on acid and base strength.

CHEMISTRY NOTES FOR CLASS - 11 (FOR PUNJAB)

Exercise

MULTIPLE CHOICE QUESTIONS (MCQs)

Q.1 Four choices are given for each question. Select the correct choice.

I. Given the following reaction:



- a) NH_3 is the acid, H_2O is the base
 b) NH_3 is the base, H_2O is the acid
 c) NH_4^+ is the base, OH^- is the acid
 d) H_2O is the base, OH^- is the acid
- II.** The pH of $10^{-3} \text{ mol dm}^{-3}$ of an aqueous solution of H_2SO_4 is:
 a) 3.0
 b) 2.7
 c) 2.0
 d) 1.5
- III.** The solubility product of AgCl is $2.0 \times 10^{-10} \text{ mol dm}^{-3}$. The maximum concentration of Ag^+ ions in the solution is:
 a) $2.0 \times 10^{-10} \text{ mol dm}^{-3}$
 b) $1.41 \times 10^{-5} \text{ mol dm}^{-3}$
 c) $1.0 \times 10^{-10} \text{ mol dm}^{-3}$
 d) $4.0 \times 10^{-20} \text{ mol dm}^{-3}$
- IV.** Which indicator is typically used for titrations involving strong acids and strong bases?
 a) Methyl red
 b) Phenolphthalein
 c) Bromothymol blue
 d) Litmus solution
- V.** Which of the following is the conjugate base of water?
 a) $\text{OH}^-(aq)$
 b) $\text{OH}^+(aq)$
 c) $\text{H}_2\text{O}(l)$
 d) $\text{H}_3\text{O}^+(aq)$
- VI.** Which of the following is a Lewis acid but not a Bronsted-Lowry acid?
 a) HCl
 b) NH_3
 c) AlCl_3
 d) H_2O
- VII.** In an acid-base titration, the equivalence point is reached when:
 a) pH of the solution is 7.0.
 b) The indicator changes color.
 c) Equal volumes of acid and base have been added.
 d) The reaction stops.
- VIII.** If the concentration of Cl^- ion in a solution is increased, the solubility of silver chloride (AgCl) will:
 a) Decrease
 b) Increase
 c) Remain unchanged
 d) Become zero
- IX.** Which of the following pairs of substances can act as a conjugate acid-base pair according to the Bronsted-Lowry theory?
 a) HCl and NaOH
 b) NH_3 and NH_4^+
 c) H_2O and H_2SO_4
 d) H_2O and CH_4
- X.** If the pH of a solution is 11, what is the $[\text{OH}^-]$ concentration in the solution?
 a) $1 \times 10^{-3} \text{ M}$
 b) $1 \times 10^{-11} \text{ M}$
 c) $1 \times 10^{-2} \text{ M}$
 d) $1 \times 10^{-14} \text{ M}$
- XI.** Which of the following pairs forms a buffer solution?
 a) HCl and NaCl
 b) CH_3COONa and CH_3COOH
 c) NaOH and HCl
 d) NH_3 and Na_2SO_4

Answer Key with Explanations

Sr.No.	Option	Answer	Explanation
I.	b	NH_3 is the base, H_2O is the acid	NH_3 accepts H^+, making it a base; H_2O donates H^+, making it an acid.
II.	b	2.7	$\text{H}_2\text{SO}_4 \rightleftharpoons 2\text{H}^+ + \text{SO}_4^{2-}$ $[\text{H}^+] = 2 \times 10^{-3}$ $\text{pH} = -\log [\text{H}^+]$ $\text{pH} = -\log 2 \times 10^{-3} = -(\log 2 - 3 \log 10) = -(0.301 - 3)$ $\text{pH} = 2.67$ Or 2.7

CHEMISTRY NOTES FOR CLASS - 11 (FOR PUNJAB)

III.	b	$1.41 \times 10^{-5} \text{ mol dm}^{-3}$	$\text{AgCl} \rightleftharpoons \text{Ag}^{+1} + \text{Cl}^{-}$ $K_{sp} = [\text{Ag}^{+1}][\text{Cl}^{-1}]$ $2.0 \times 10^{-10} = [\text{Ag}^{+1}][\text{Cl}^{-1}]$ Since $[\text{Ag}^{+1}] = [\text{Cl}^{-1}]$ $2.0 \times 10^{-10} = [\text{Ag}^{+1}]^2$ $\sqrt{2.0 \times 10^{-10}} = \sqrt{[\text{Ag}^{+1}]^2}$ $[\text{Ag}^{+1}] = 1.414 \times 10^{-5} \text{ mol dm}^{-3}$
IV.	b	Phenolphthalein	Changes color near pH 7–9, ideal for strong acid-strong base titrations.
V.	a	$\text{OH}^{-}_{(\text{aq})}$	H_2O loses H^{+} to form OH^{-} so OH^{-} is its conjugate base.
VI.	c	AlCl_3	Accepts an electron pair but does not donate H^{+}.
VII.	c	Equal volumes of acid and base have been added.	Equivalence point is stoichiometric, not necessarily pH 7.
VIII.	a	Decrease	Common ion effect reduces solubility.
IX.	b	NH_3 and NH_4^{+}	NH_3 gains H^{+} to form NH_4^{+}
X.	a	$1 \times 10^{-3} \text{ M}$	$\text{pH} = 11 \rightarrow \text{pOH} = 3 \rightarrow [\text{OH}^{-}] = 10^{-3} \text{ M}$.
XI.	b	Sodium acetate (CH_3COONa) and acetic acid (CH_3COOH)	Weak acid (CH_3COOH) + its conjugate base ($\text{CH}_3\text{COO}^{-}$) forms buffer.

SHORT ANSWER QUESTIONS

Q.2 Attempt the following short-answer questions:

a. Define the following with an example for each:

i) Ionization constant

ii) Solubility product


iii) Common ion effect

iv) Acid-base Indicator

Ans.


(i) **Ionization Constant:** The quantitative measure of the strength of the acid and base is called ionization constant.

Example: we have an acid HA dissolved in water,





(ii) **Solubility Product:** The solubility product is the product of the concentrations of ions raised to an exponent equal to the co-efficient of the balanced equation.

Example: For AgCl



(iii) **Common ion Effect:** The suppression of ionization of a weak electrolyte by adding a common ion to it is called common ion effect.



(iv) **Acid-base Indicator:** An indicator is a substance that changes colour to mark a titration's endpoint. Acid-base indicators exhibit one colour in acid and another in base. Most indicators used in acid-base titration are weak organic acids or weak organic bases.

Example:

Phenolphthalein: Colorless (pH < 8.3) to Pink (pH > 10.0)

CHEMISTRY NOTES FOR CLASS - 11 (FOR PUNJAB)

b. Differentiate between:

i) Hydrolysis and dissolution

ii) Acidic and basic buffer solutions

Ans. i) Hydrolysis and Dissolution

Feature	Hydrolysis	Dissolution
Definition	A chemical reaction where water breaks bonds in a compound.	A physical process where a solute dissolves in a solvent.
Nature	Chemical process (forms new substances).	Physical process (no new substances formed).
Involves	Reaction with water molecules (H^+ and OH^- may participate).	Mixing of solute and solvent (e.g., salt in water).
Energy Change	May release or absorb energy (exo/endothermic).	Often absorbs energy (endothermic, but not always).
Result	Forms new products (e.g., acids, bases, or ions).	Forms a homogeneous mixture (solution).
Example	$AlCl_3 + 3H_2O \rightleftharpoons Al(OH)_3 + 3H^+ + 3Cl^-$	$NaCl_{(s)} \rightarrow Na^+_{(aq)} + Cl^-_{(aq)}$

(ii) Acidic and Basic Buffer Solutions

Acidic buffers	Basic buffers
- Those buffers which are formed by the mixing of a weak acid and a salt of it with a strong base are called acidic buffers. - Their pH is less than 7. - Henderson's equation for acidic buffers $pH = pK_a + \log \frac{[salt]}{[acid]}$	- Those buffers which are formed by mixing of a weak base and a salt of it with a strong acid are called basic buffers. - Their pH is more than 7. - Henderson's equation for basic buffers $pOH = pK_b + \log \frac{[salt]}{[base]}$
e.g. $CH_3COOH + CH_3COONa$ weak acid salt of weak acid	e.g. $NH_4OH + NH_4Cl$ weak base salt of weak base

c. Explain the concept of conjugate acid-base pairs. How are they related in terms of proton transfer?


Ans. Conjugate Acid:

"A species formed after a Bronsted base accepts a proton from the acid is called the conjugate acid". The hydronium ion (H_3O^+) is the conjugate acid of water.

Example:

When NH_3 dissolves in Water:

When ammonia dissolves in water, proton transfer also occurs. H^+ from H_2O attaches to the N atom's lone pair and NH_4^+ and OH^- are formed. With fewer H^+ , the water molecule becomes OH^- ion:




In this case, H_2O (the acid) has donated the proton and NH_3 (the base) has accepted it. $NH_3 - NH_4^+$ and $H_2O - OH^-$ are conjugate acid-base pair.

Conjugate Base:

"A species formed after a Bronsted base accepts a proton from the acid is called the conjugate acid."

Example: The hydronium ion (H_3O^+) is the conjugate acid of water. A species formed when an acid donates a proton to a base is called the conjugate base. $HCl + Cl^-$ and $H_2O \rightarrow H_3O^+$ are conjugate acid-base pairs.



CHEMISTRY NOTES FOR CLASS - 11 (FOR PUNJAB)

d. What is the relationship between the strength of an acid and the strength of its conjugate base?


Ans. A strong acid has a weak conjugate base because it donates protons easily. A weak acid has a stronger conjugate base as it holds onto protons.

Example: HCl (strong acid) forms Cl^- (very weak base).

- Strong Acid (HCl): Weak conjugate base (Cl^-).
- Weak Acid (CH_3COOH): Strong conjugate base (CH_3COO^-).

21
21

e. For the following three reactions, identify the reactants that are Arrhenius bases, Bronsted-Lowry bases, and/or Lewis bases. State which type(s) of bases each reactant is. Explain your answers.




Ans. (i) OH^- is an Arrhenius and Bronsted-Lowry base (accepts H^+).

(ii) H_2O is a Bronsted-Lowry base (accepts H^+) and Lewis base (donates lone pair).

(iii) NH_3 is a Bronsted-Lowry and Lewis base (accepts H^+ and donates lone pair).

f. An amphoteric substance can behave as either an acid or a base. Identify whether water behaves as an acid or a base in each of the following reactions.



Ans.	Reaction	H_2O Role	Reason
i)	$\text{H}_2\text{O} + \text{HCl} \longrightarrow \text{H}_3\text{O}^+ + \text{Cl}^-$	Base	Accepts H^+ from HCl .
ii)	$\text{NH}_3 + \text{H}_2\text{O} \longrightarrow \text{NH}_4^+ + \text{OH}^-$	Acid	Donates H^+ to NH_3 .
iii)	$\text{HNO}_3 + \text{H}_2\text{O} \longrightarrow \text{H}_3\text{O}^+ + \text{NO}_3^-$	Base	Accepts H^+ from HNO_3 .
iv)	$\text{CH}_3\text{COOH} + \text{H}_2\text{O} \longrightarrow \text{CH}_3\text{COO}^- + \text{H}_3\text{O}^+$	Base	Accepts H^+ from CH_3COOH .

g. Which salt would you expect to dissolve more readily in acidic solution: Barium sulfate or Barium fluoride? Explain. $K_{sp}(\text{BaCO}_3) = 1.1 \times 10^{-10}$, $K_{sp}(\text{CuS}) = 8.1 \times 10^{-36}$

Ans. Barium fluoride (BaF_2) dissolves more in acid.


Reason: F^- reacts with H^+ to form HF :



BaSO_4 is unaffected because SO_4^{2-} doesn't react with H^+ .

h. Why does common ion effect decrease solubility of a less soluble salt?

Ans. The solubility of a less soluble salts KClO_3 in water is suppressed by the addition of a more soluble salt KCl by common ion effect. K^+ is a common ion. The ionization of KClO_3 is suppressed and it settles down as precipitate.



i. State the basic principle of solubility product. Mention factors affecting solubility product.

Ans. It is the product of molar solubilities of two ions at equilibrium stage.

For example: sparingly soluble salt, PbCl_2 . $K_{sp} = [\text{Pb}^{2+}][\text{Cl}^-]^2$

K_{sp} is usually a very small quantity at room temperature. The value of K_{sp} is temperature dependent.


Factors Affecting K_{sp} :

- Temperature
- Presence of common ions
- pH (if salt has basic/acidic ions)

CHEMISTRY NOTES FOR CLASS - 11 (FOR PUNJAB)

j. Prove by equations what happens when Na_2CrO_4 is added to saturated solution of PbCrO_4 .

Ans. When Na_2CrO_4 is added to a saturated PbCrO_4 solution, it increases the concentration of CrO_4^{2-} ions. According to Le Chatelier's Principle, the equilibrium shifts left to reduce the added CrO_4^{2-} ions:



- **Result:** More PbCrO_4 precipitates out, decreasing its solubility.
- **Reason:** The common ion (CrO_4^{2-}) suppresses the dissociation of PbCrO_4 .
This is an example of the common ion effect reducing solubility.

k. According to the Lewis acid-base concept, Boron trifluoride (BF_3) can act as an acid. Is this statement correct?

Ans. Correct, BF_3 is a Lewis acid.

Reason: Boron in BF_3 has an empty p-orbital and can accept an electron pair (Lewis acid).

Example: When BF_3 reacts with NH_3 (a Lewis base), it forms a coordinate covalent bond:



Here, BF_3 accepts the lone pair from NH_3 .

l. If the concentration of hydrogen ions in a solution is $1 \times 10^{-5} \text{ M}$, what is the pH of the solution?

Given Data:

Concentration of hydrogen ions, $[\text{H}^+] = 1 \times 10^{-5} \text{ M}$

To Find:

pH of the solution = ?

Solution:

Step 1: pH formula:

$$\text{pH} = -\log[\text{H}^+]$$

Step 2: Substitute the given $[\text{H}^+]$ into the formula:

$$\text{pH} = -\log(1 \times 10^{-5})$$

Step 3: Simplify the logarithm:

$$\text{pH} = -(-5) = 5$$

Answer: The pH of the solution is 5.

DESCRIPTIVE QUESTIONS

Q.3 Describe the Bronsted-Lowry theory of acids and bases. Provide examples of conjugate acid-base pairs and explain clearly their relationship.

Ans. See Page No. (294)

Q.4 Define the Lewis theory of acids and bases. How does this theory differ from the Bronsted-Lowry theory? Give examples of Lewis acids and bases that do not involve proton transfer.

Ans. See Page No. (295)

Q.5 Discuss applications and implications of the common ion effect in various fields.

Ans. See Page No. (305)

Q.6 What is the solubility product for sparingly soluble salts? Give its two applications.

Ans. See Page No. (309)

Q.7 Describe the general shape of a titration curve for a strong acid titrated with a strong base. How can you identify the equivalence point on this titration curve?

Ans. See Page No. (315)

CHEMISTRY NOTES FOR CLASS - 11 (FOR PUNJAB)

NUMERICAL PROBLEMS

Q.8 A buffer solution has a pH of 5.0. It is made from a weak acid HA with a pK_a of 4.8. What is the ratio of the concentration of the conjugate base $[A^-]$ to the concentration of the weak acid $[HA]$ in this buffer?

Given Data:

$$pH = 5.0$$

$$pK_a = 4.8$$

Formula:

$$pH = pK_a + \log \frac{[salt]}{[acid]}$$

Putting the values:

$$5.0 = 4.8 + \log \left(\frac{[A^-]}{[HA]} \right)$$

Solve for the log term: $\log \left(\frac{[A^-]}{[HA]} \right) = 5.0 - 4.8 = 0.2$

Convert log to ratio (antilog):

$$\frac{[A^-]}{[HA]} = 10^{0.2} \approx 1.58$$

Final Result:

$$\frac{[A^-]}{[HA]} \approx \boxed{1.58}$$

The buffer contains 1.58 times more conjugate base (A^-) than weak acid (HA) to maintain pH 5.0.

Q.9 Calculate the solubility of a sparingly salt lead (II) iodide (PbI_2) in water. It has $K_{sp} = 1.4 \times 10^{-8}$.

Given Data:

$$K_{sp} = 1.4 \times 10^{-8}$$



Define Solubility (s):

Let

At equilibrium:

$$s = \text{molar solubility of } PbI_2 \text{ (mol/dm}^3\text{)}.$$

$$[Pb^{2+}] = s$$

$$[I^-] = 2s \text{ (2 moles of } I^- \text{ are produced per mole of } PbI_2\text{).}$$

K_{sp} Expression:

$$K_{sp} = [Pb^{2+}][I^-]^2$$

$$= (s)(2s)^2 = 4s^3$$

Solve for s:

$$4s^3 = 1.4 \times 10^{-8}$$

$$s^3 = \frac{1.4 \times 10^{-8}}{4} = 3.5 \times 10^{-9}$$

$$s = \sqrt[3]{3.5 \times 10^{-9}} \approx \boxed{1.52 \times 10^{-3} \text{ mol/dm}^3}$$

Final Answer: The solubility of PbI_2 is $1.52 \times 10^{-3} \text{ mol/dm}^3$.

Q.10 The molar solubility of silver chromate (Ag_2CrO_4) in pure water at 298 K is $6.5 \times 10^{-5} \text{ mol dm}^{-3}$. Calculate the K_{sp} of silver chromate at this temperature.

Given Data:

$$\text{Molar solubility (s) of } Ag_2CrO_4 = 6.5 \times 10^{-5} \text{ mol dm}^{-3}$$



Ion Concentrations in Terms of s

$$[Ag^+] = 2s \text{ (2 moles of } Ag^+ \text{ per mole of } Ag_2CrO_4\text{).}$$

$$[CrO_4^{2-}] = s \text{ (1 mole of } CrO_4^{2-} \text{ per mole of } Ag_2CrO_4\text{).}$$

K_{sp} Expression:

$$K_{sp} = [Ag^+]^2[CrO_4^{2-}]$$

$$= (2s)^2(s) = 4s^3$$

Substitute s and Solve for K_{sp} :

$$K_{sp} = 4 \times (6.5 \times 10^{-5})^3$$

$$K_{sp} = 4 \times (2.746 \times 10^{-13})$$

$$K_{sp} = 1.10 \times 10^{-12} \text{ mol}^3 \text{ dm}^{-9}$$

Final Answer:

$$K_{sp} = \boxed{1.1 \times 10^{-12}}$$

CHEMISTRY NOTES FOR CLASS - 11 (FOR PUNJAB)

ADDITIONAL SLOs BASED MCQs

- Water cannot act as:
 - Lewis acid
 - Lewis base
 - Bronsted acid
 - Bronsted base
- pH of 0.01 M HCl solution is:
 - 10^{-2}
 - 10^{+2}
 - 2.0
 - 1.0
- An aqueous solution of ammonium chloride is:
 - Basic
 - Acidic
 - Neutral
 - Amphoteric
- An aqueous solution of which compound is basic.
 - Ammonium nitrate
 - Calcium chloride
 - Ammonium acetate
 - Potassium carbonate
- Which statement about acids is not correct? An acid:
 - Contains hydrogen ions in solution
 - Contains Oxygen
 - Has a pH of less than 7 then
 - Gives off carbon dioxide from a carbonate
- If a liquid has a pH of 7 then:
 - It must be colourless
 - It has boiling point of 100°C
 - It must be a solution
 - It must be neutral
- When air is bubbled through pure water, the pH is lowered from 7.0 to 5.6, which gas in the air is responsible for this change.
 - Argon
 - Carbon dioxide
 - Nitrogen
 - Oxygen
- Which of the following oxides is classified incorrectly?
 - Zinc oxide (ZnO) amphoteric
 - Carbon dioxide (CO_2) acidic
 - Carbon monoxide (CO) neutral
 - Lead oxide (PbO) basic
- If 25cm^3 of 1 mole. dm^{-3} nitric acid is added to 50cm^3 of 0.5 potassium hydroxide solution, what would be the pH of the resulting solution?
 - 5
 - 7
 - 9
 - 14
- If dry citric acid crystals are placed on dry litmus paper they will:
 - turn yellow
 - turn green
 - turn red
 - remain unchanged
- A base is a substance which will neutralize an acid. Which of these substances is not a base?
 - Aqueous ammonia
 - Copper oxide
 - Potassium chloride
 - Sodium carbonate
- A strong acid:
 - Is always partially ionized when in solution
 - Is always fully ionized when in solution
 - Always decomposes carbonates
 - Always contains oxygen
- Which one of the following oxides dissolves in water to form acidic solution?
 - MgO
 - Na_2O
 - SO_2
 - SiO_2
- When crystals of copper sulphate are heated, the colour changes from blue to white. This is caused by:
 - Loss of water only
 - Loss of water and SO_2
 - Reaction with CO_2 in the air
 - Loss of water, sulphur dioxide and oxygen
- The oxide of a metal was found to react with HCl and aqueous NaOH solution. Which of the following is the best description of the oxide?
 - Acidic
 - Amphoteric
 - Basic
 - Neutral

CHEMISTRY NOTES FOR CLASS - 11 (FOR PUNJAB)

Answer Key with Explanations

Q. No	Ans.	Explanation												
1.	A	- Water can donate and accept proton (H^+). So it can act as Bronsted acid as well as base. $H_2O + H_2O \rightleftharpoons H_3O^+ + OH^-$ - Similarly, in water, oxygen has electron pairs which it can donate to other species, so water acts as Lewis base. $H-\ddot{O}: + H^+ \rightarrow \left[H-\ddot{O} \rightarrow H \right]^+$ But water cannot act as Lewis acid.												
2.	C	$HCl \rightleftharpoons H^+ + Cl^-$ 0.01M 0.01M Since HCl is a strong acid, it ionized completely giving $[H^+] = 0.01M = 10^{-2}M$ $pH = -\log [H^+]$ $pH = -\log 10^{-2}$ $pH = 2 \log 10 \quad \log 10 = 1$ $pH = 2$												
3.	B	$HCl + NH_3 \rightleftharpoons NH_4Cl$ Strong acid weak base Acidic salt When NH_4Cl is added, the solution becomes acidic.												
4.	D	<table border="1"> <thead> <tr> <th>Salt</th><th>Type</th><th>Nature of Solution</th></tr> </thead> <tbody> <tr> <td>NH_4NO_3 Ammonium Nitrate</td><td>Acidic Salt</td><td>Acidic solution</td></tr> <tr> <td>$CaCl_2$ Calcium chloride</td><td>Neutral salt</td><td>Somewhat neutral solution</td></tr> <tr> <td>K_2CO_3 Potassium carbonate</td><td>Basic salt</td><td>Basic solution</td></tr> </tbody> </table>	Salt	Type	Nature of Solution	NH_4NO_3 Ammonium Nitrate	Acidic Salt	Acidic solution	$CaCl_2$ Calcium chloride	Neutral salt	Somewhat neutral solution	K_2CO_3 Potassium carbonate	Basic salt	Basic solution
Salt	Type	Nature of Solution												
NH_4NO_3 Ammonium Nitrate	Acidic Salt	Acidic solution												
$CaCl_2$ Calcium chloride	Neutral salt	Somewhat neutral solution												
K_2CO_3 Potassium carbonate	Basic salt	Basic solution												
5.	B	$HCl, HBr, BF_3, AlCl_3$ etc are acids but they do not contain oxygen. Anyhow H_2SO_4, HNO_3, CH_3COOH etc have oxygen in them												
6.	D	Solution $\begin{cases} pH = 7 \text{ (Neutral)} \\ pH > 7 \text{ (Basic)} \\ pH < 7 \text{ (Acidic)} \end{cases}$												
7.	B	When air is bubbled through pure water, CO_2 combines with water and forms carbonic acid. $CO_{2(g)} + H_2O_{(l)} \rightarrow H_2CO_{3(aq)}$ As a result, pH of the solution decreases												
8.	D	Lead oxide (PbO) is an amphoteric oxide. $PbO + H_2SO_4 \rightarrow PbSO_4 + H_2O$ (base) (acid) $PbO + 2NaOH + H_2O \rightarrow Na_2[Pb(OH)_4]$ (acid) (base)												
9.	B	Concentration = Molarity $\times$ Volume of H^+ or OH^- Since $[H^+] = [OH^-]$ in both cases, so a neutral solution having $pH = 7$ is formed.												
10.	D	Moisture is necessary to effect the litmus paper. In moist, the litmus paper turns red with citric acid.												
11.	C	$KOH + HCl \rightarrow KCl + H_2O$ Strong base Strong acid Neutral salt												


CHEMISTRY NOTES FOR CLASS - 11 (FOR PUNJAB)

		KCl will not hydrolyse, so solution remains neutral.
12.	B	The strength of an acid depends upon its extent of ionization. A strong acid fully ionizes in solution.
13.	C	Generally – Metals form basic oxides Non-metals form acidic oxides $\text{SO}_2 + \text{H}_2\text{O} \longrightarrow \text{H}_2\text{SO}_3 \text{ (sulphurous acid)}$ Acidic oxide
14.	A	Blue vitriol is deep blue solid copper (II) sulphate pentahydrate is heated to drive off water vapours to form white solid copper (II) sulphate.
15.	B	Aluminum oxide (Al_2O_3) is amphoteric and it reacts with both acidic and basic solution. $\text{Al}_2\text{O}_3(s) + 6\text{HCl}(aq) \longrightarrow 2\text{AlCl}_3(aq) + 3\text{H}_2\text{O}(l)$ (basic) $\text{Al}_2\text{O}_3(s) + 2\text{NaOH}(aq) \longrightarrow 2\text{NaAlO}_2(aq) + \text{H}_2\text{O}(l)$ (acidic)


ADDITIONAL SHORT ANSWER QUESTIONS

Q.1 Why do the equilibrium constant value has its units for some of the reversible reactions but has no units for some other reactions?

Ans. i. If number of moles of reactants and products are equal, then K_c has no units.



ii. If the number of moles of reactants and products are not equal.



Q.2 Define pH and pOH with mathematical expressions. What is the sum of pH and pOH?

Ans. pH: "The negative log of hydrogen ion concentration is called pH."

$$\text{pH} = -\log[\text{H}^+]$$

pOH: "The negative log of concentration of hydroxide ion is called pOH."

$$\text{pOH} = -\log[\text{OH}^-]$$

Relationship between pH and pOH:

$$\text{pH} + \text{pOH} = 14 \text{ at } 25^\circ\text{C}$$

The sum of pH and pOH for a solution must be equal to 14 at 25°C .

Q.3 Write the relationship of pH and pOH with pK_w .

Ans. Relation of pH and pOH with pK_w :

From the concept of ionic product of water we have

$$K_w = [\text{H}^+][\text{OH}^-]$$

Taking log:

$$\log K_w = \log[\text{H}^+] + \log[\text{OH}^-]$$

Now multiply with (-1)

$$-\log K_w = -\log[\text{H}^+] + (-\log[\text{OH}^-])$$

$$\text{pK}_w = \text{pH} + \text{pOH}$$

$$\text{pH} + \text{pOH} = 14 \text{ at } 25^\circ\text{C} \quad (\text{Since, } \text{pK}_w = 14 \text{ at } 25^\circ\text{C})$$

Thus, the sum of pH and pOH is equal to pK_w i.e., 14 at 25°C .

CHEMISTRY NOTES FOR CLASS - 11 (FOR PUNJAB)

Q.4 What is ionization constant of acids.


Ans. Ionization Constant of Acids:

The ionization constants of the acids is the quantitative measure of the strength of the acid. Acids and bases when dissolved in water may or may not be completely dissociated. Many acids are weak electrolytes and they ionize to an extent which is less than 100%.

Q.5 Define Lowry Bronsted concept of acids and bases.

Ans. Bronsted Acid:

"Those species (ions or molecules) which donate or have a tendency to donate proton [H⁺] are called acids."



Bronsted Base:

"These species (ions or molecules) which accept or have a tendency to accept proton (H⁺) are called bases."

Q.6 What are conjugate acids and bases?


Ans. Conjugate Acid:

A conjugate acid is the specie (molecule or ion) formed when a proton is added to a base. e.g., When a weak acid or a weak base is dissolved in water, the conjugate acid base pair is produced.



Conjugate base: The species (ions or molecules) formed when acid loses its H⁺ ions and resulting species become its conjugate base.


Example:



Conjugate acid-base Pair

When a weak acid or a weak base is dissolved in water, the conjugate acid base pair is produced.

Example:



Q.7 What is the formula to calculate the percentage ionization of weak acid? (LHR 2018 GII)

Ans. Percentage of Ionization of Weak Acids:

We can calculate the percentage of ionization of weak acid and the formula is as follows:

$$\% \text{ Ionization} = \frac{\text{Amount of acid ionized}}{\text{Amount of acid initially available}} \times 100$$

Q.8. Define pK_a and pK_b. Give its significance.

Ans. pK_a:

"The negative log of dissociation constant of acid (K_a) is called pK_a".

$$pK_a = -\log K_a$$

$$pK_a \propto \frac{1}{K_a}$$

pK_b:

"The negative log of K_b is called pK_b".

$$pK_b = -\log K_b$$

CHEMISTRY NOTES FOR CLASS - 11 (FOR PUNJAB)

Significance:

- pK_a and pK_b gives us the quantitative strength of acids and bases respectively.
- Larger the value of pK_a , weaker will be the acid and vice versa.
- Larger the value of pK_b , weaker will be the base and vice versa.

Q.9 Prove that $pK_a + pK_b = 14$ at 25°C .

Ans. Let us take the log of equation:

$$\log(K_a \times K_b) = \log(K_w)$$

$$\log K_a + (\log K_b) = \log K_w$$

Multiply both sides by '-1'

$$-\log K_a - \log K_b = -\log K_w$$

Since, $pK_a = -\log K_a$ and $pK_b = -\log K_b$

So, $pK_a + pK_b = pK_w = 14$

Hence pK_a and pK_b of conjugate acid base pair has a very simple relation with each other.

$$pK_a + pK_b = 14 \text{ at } 25^\circ\text{C}$$

Q.10 What is ionic product of water? Write its value at 25°C .

Ans. Ionic Product of Water:

The product of molar concentration of H^+ and OH^- in pure water is called ionic product of water. It is represented by K_w and its value is 10^{-14} at 25°C .

$$K_w = [\text{H}^+][\text{OH}^-]$$

$$K_w = 10^{-14} \text{ at } 25^\circ\text{C}$$

Q.11 What is buffer solution? Give types of buffer solution with one example each.

Ans. Buffer Solutions:

"These solutions which resist the change in their pH when a small amount of an acid or a base is added to them are called Buffer solution."

The buffer solutions have a specific constant value of pH.

The pH values of buffer solutions do not change on dilution and on keeping for a long time.

Types of Buffer Solutions: Buffer solutions are mostly prepared by mixing two substances.

Acidic Buffers	Basic Buffers
• These buffers which are formed by the mixing of a weak acid and a salt of it with a strong base are called acidic buffers. • Their pH is usually less than 7. • Henderson's equation for acidic buffers $\text{pH} = \text{pK}_a + \log \frac{[\text{salt}]}{[\text{acid}]}$	• These buffers which are formed by mixing of a weak base and a salt of it with a strong acid are called basic buffers. • Their pH is more than 7. • Henderson's equation for basic buffers $\text{pOH} = \text{pK}_b + \log \frac{[\text{salt}]}{[\text{base}]}$
e.g., CH_3COOH + CH_3COONa weak acid salt of weak acid	e.g., NH_4OH + NH_4Cl weak base salt of weak base

Q.12 Define buffer capacity. Write down the Henderson's equation for acidic buffers.

Ans. Buffer Capacity:

"The capability of a buffer solution to resist the change in its pH when a small amount of acid or a base is added in it is called buffer capacity."

$$\text{pH} = \text{pK}_a + \log \frac{[\text{Salt}]}{[\text{Acid}]}$$

CHEMISTRY NOTES FOR CLASS - 11 (FOR PUNJAB)


Q.13 Define common ion effect. How can NaCl be purified by common ion effect?

Ans. Common Ion Effect:

"The suppression of ionization of a weak electrolyte by adding a common ion from outside is called common ion effect."

Example: Purification of NaCl:

NaCl is purified by passing hydrogen chloride gas through the saturated brine. Sodium chloride is fully ionized in the solution. Equilibrium constant expression for this process can be written as follows:



HCl also ionizes in solution.



On passing HCl gas, concentration of Cl^- ions is increased. Therefore, NaCl crystallizes out of the solution to maintain the constant value of equilibrium constant. This type of effect is called common ion effect.



The addition of a common ion to the solution of less soluble electrolyte suppresses its ionization and the concentration of unionized species increases, which may come out as a precipitate.

Q.14 Give applications of common ion effect (any two).

Ans. Applications of Common Ion Effect:

- **Precipitation of 2nd Group Basic Radicals of Salt Analysis:**

Similarly the dissociation of a weak acid H_2S in water can be suppressed by the addition of stronger acid HCl. H^+ is a common ion. H_2S becomes less dissociated in acidic solution. In this way, low concentration of S^{2-} is developed.




This low concentration of S^{2-} ions helps to do the precipitation of radicals of second group basic radicals during salt analysis.



- **Precipitation of 3rd Group Basic Radicals of Salt Analysis:**


An addition of NH_4Cl in NH_4OH solution suppresses the concentration of $\text{OH}^-_{(aq)}$ due to the presence of a large excess of NH_4^+ from NH_4Cl . Actually, NH_4Cl is a strong electrolyte. The combination of these two substances is used as reagent in third group basic radicals.



Q.15 Write the effect of common ion on solubility. Prove by equations that what happens when Na_2CrO_4 is added to saturated solution of PbCrO_4 ?

Ans. Effect of Common Ion on Solubility:

The presence of a common ion decreases the solubility of a slightly soluble ionic compound. In order to explain it, consider a saturated solution of PbCrO_4 , which is a sparingly soluble ionic salt.



Now add Na_2CrO_4 , which is a soluble salt. CrO_4^{2-} is the common ion. It combines with Pb^{2+} to form more insoluble PbCrO_4 . So equilibrium is shifted to the left to keep K_{sp} constant.

CHEMISTRY NOTES FOR CLASS - 11 (FOR PUNJAB)

Q.16 Define Solubility Product Constant (K_{sp}). Give two examples.

Ans. Solubility Product:

“The product of the equilibrium concentration of ions raised to an exponent equal to the coefficient of the balanced equation at given temperature is called solubility product (K_{sp}).”

Mathematical Form:



$$K_c = \frac{[A^{+y}]^x [B^{-x}]^y}{[A_xB_y]}$$


$$K_c [A_xB_y] = [A^{+y}]^x [B^{-x}]^y$$

$$K_{sp} = [A^{+y}]^x [B^{-x}]^y$$

- K_{sp} is the product of molar solubilities of two ions at equilibrium.

Examples:

1. When Lead Chloride ($PbCl_2$) is shaken with water the solution contains Pb^{+2} , Cl^- and un-dissociated $PbCl_2$. It means an equilibrium exists between $PbCl_2$ (solid solute) and the dissolved ions, Pb^{+2} and Cl^- .



According to law of mass action:

$$K_c = \frac{[Pb^{+2}][Cl^-]^2}{[PbCl_2]}$$

$$K_c [PbCl_2] = [Pb^{+2}][Cl^-]^2$$

$$K_{sp} = [Pb^{+2}][Cl^-]^2 = 1.6 \times 10^{-5} \text{ at } 25^\circ\text{C}$$

2. **Solubility Product Expression for Ag_2CrO_4 :**



$$K_c = \frac{[Ag^+]^2 [CrO_4^{2-}]}{[Ag_2CrO_4]}$$

$$K_{sp} = K_c [Ag^+]^2 [CrO_4]^{-2}$$



CHEMISTRY NOTES FOR CLASS - 11 (FOR PUNJAB)

SELF-ASSESSMENT Chapter # 09

Total Mark: 30

Q.1 Encircle the correct option.

(1 × 6 = 6)

(i) Given the following reaction:



- A. NH_3 is the acid, H_2O is the base
 B. NH_3 is the base, H_2O is the acid
 C. NH_4^+ is the base, OH^- is the acid
 D. H_2O is the base, OH^- is the acid

(ii) In the given reaction, the conjugate base is:



- A. NH_3 B. Cl^- C. NH_4^+ D. HCl

(iii) Which statement correctly describes a Lewis acid?

- A. Accepts a lone pair of electrons
 B. Donates a lone pair of electrons
 C. Always donates protons
 D. Has a negative charge

(iv) If the pH of a solution is 11, what is the $[\text{OH}^-]$ concentration in the solution?

- A. $1 \times 10^{-3} \text{ M}$ B. $1 \times 10^{-11} \text{ M}$ C. $1 \times 10^{-2} \text{ M}$ D. $1 \times 10^{-14} \text{ M}$

(v) The solubility product of AgCl is $2.0 \times 10^{-10} \text{ mol dm}^{-3}$. The maximum concentration of Ag^+ ions in the solution is:

- A. $2.0 \times 10^{-10} \text{ mol dm}^{-3}$ B. $1.41 \times 10^{-5} \text{ mol dm}^{-3}$ C. $1 \times 10^{-10} \text{ mol dm}^{-3}$ D. $4.0 \times 10^{-20} \text{ mol dm}^{-3}$

(vi) Which indicator is typically used for titrations involving strong acids and strong bases?

- A. Methyl red B. Phenolphthalein C. Bromothymol blue D. Litmus solution

Q.2 Write short answers of the following questions.

(2 × 8 = 16)

(i) Identify the Lewis acid and Lewis base in the reaction between:

- (a) SO_3 and O^{2-} (b) HCl and H_2O

Also write down the balanced chemical equation for the reaction and explain

(ii) Calculate the pH of a 0.01 M NaOH solution.

(iii) Determine $[\text{H}_3\text{O}^+]$ in a 0.01 M HA solution given $K_a = 1.0 \times 10^{-6}$.

(iv) Define common ion effect. Give a suitable example.

(v) Differentiate end point and equivalent point.

(vi) A solution is prepared by mixing equal volumes of two solutions: one with a pH of 4.0 and another with a pH of 10.0. Calculate the K_w for this mixture at 25°C.

(vii) How does a buffer maintain pH stability?

(viii) Why is aqueous solution of NH_4Cl slightly acidic?

Q.3 Extensive Questions.

(2 × 4 = 8)

(a) Describe the Bronsted-Lowry theory of acids and bases. Provide examples of conjugate acid-base pairs and explain clearly their relationship.

(b) The molar solubility of silver chromate (Ag_2CrO_4) in pure water at 298 K is $6.5 \times 10^{-5} \text{ mol dm}^{-3}$. Calculate the K_{sp} of silver chromate at this temperature.

