

UNIT

10

INTRODUCTION TO COORDINATE GEOMETRY

- ▶ Introduction To Coordinate Geometry
- ▶ Distance Formula
- ▶ Collinear Points

After completion of this unit, the students will be able to:

- ▶ define coordinate geometry.
- ▶ derive distance formula to find distance between two points given in cartesian plane.
- ▶ use distance formula to find distance between two given points.
- ▶ define collinear points.
- ▶ distinguish between collinear and non-collinear points.
- ▶ use distance formula to show that given three (or more) points are collinear.
- ▶ use distance formula to show that the given three non-collinear points form:
 - An equilateral triangle.
 - An isosceles triangle.
 - A right angled triangle.
 - A scalene triangle.

10.1 DISTANCE FORMULA

In 17th Century, Descartes, a French Mathematician introduced a plane. A set of infinite number of points, called Cartesian Plane. Every point in a plane can be located in terms of a pair of numbers related to two number axes in the plane, which are perpendicular to each other and intersect at the origin.

The plane is called a cartesian plane, and the axes are designated the horizontal $(\overline{OX})$ and vertical $(\overline{OY})$ axes, or the x -axis and the y -axis.



axes divide the plane into four quadrants shown in the figure.

in the plane and a line through P is perpendicular to the x -axis, the lines will intersect at two points,

coordinate of $M(m)$

on the x -axis is called the x -coordinate or abscissa of P , and the coordinate of $N(n)$ on the y -axis is called the y -coordinate or ordinate of P .

The two numbers (m,n) are called the coordinates of P with respect to the coordinate axes. The letters m and n stand for numbers, and since the x -coordinate is always written first, such a pair is called an ordered pair of numbers. That is, the pair $(3,2)$ is not the same as the pair $(2,3)$.



Remember that:

- (i) A point in a number plane determines a unique ordered pair of numbers.
- (ii) With every ordered pair of numbers a unique point is associated in the plane.

Since numbers to the right of the origin on the horizontal axis and numbers above the origin on the vertical axis are taken as positive, therefore:


- (i) A point in the *Ist* quadrant is characterized by the fact that both its coordinates are positive.
- (ii) A point in the *IInd* quadrant has its abscissa negative and its ordinate positive.
- (iii) A point in the *IIIrd* quadrant has both coordinates negative.
- (iv) A point in the *IVth* quadrant has its abscissa positive and its ordinate negative.
- (v) Points on the axes do not lie in any quadrant.
- (vi) Points on the positive x -axis have a positive abscissa, and their ordinate is "0".
- (vii) Points on the negative x -axis have a negative abscissa, and their ordinate is "0".
- (viii) Points on the positive y -axis have a positive ordinate, and abscissa is "0".
- (ix) Points on the negative y -axis have a negative ordinate, and their abscissa is "0".
- (x) The origin has the coordinates $(0, 0)$.

EXAMPLE-1


Locate $(2, -4)$ in the co-ordinate plane.

SOLUTION:

In this problem abscissa is positive, therefore it would be towards the right of the origin, and the ordinate is negative, so it would be below the origin, therefore the given point P is as shown in the figure.

**10.1.2 Distance Between Two Points**

Consider the points $P(x_1, y_1)$ and $Q(x_2, y_2)$ in the cartesian plane as shown in the figure. To find the length of the segment $\overline{PQ}$, we form a right triangle by drawing through P a line parallel to x -axis and through Q a line parallel to y -axis and let these lines meet at $R(x_2, y_1)$.



By Pythagoras theorem, we have.

$$|\overline{PQ}|^2 = |\overline{PR}|^2 + |\overline{RQ}|^2$$

$$|\overline{PQ}|^2 = |(x_2 - x_1)|^2 + |(y_2 - y_1)|^2$$

$$= (x_2 - x_1)^2 + (y_2 - y_1)^2$$

$$\text{Hence } |\overline{PQ}| = \pm \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

As we are only interested in the length of the segment and not in the direction, therefore we only consider the positive sign.

Hence distance between two points $P(x_1, y_1)$ and $Q(x_2, y_2)$ is given by:

$$d = |\overline{PQ}| = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

10.1.3 Use of Distance Formula

EXAMPLE-1

What kind of a triangle has vertices $A(6, -2)$, $B(1, -2)$ and $C(-2, 2)$?

SOLUTION:

Given $A(6, -2)$, $B(1, -2)$ and $C(-2, 2)$. Using distance formula,

$$|\overline{AB}| = \sqrt{(1-6)^2 + (-2+2)^2} = \sqrt{5^2 + 0} = \sqrt{25} = 5$$

$$|\overline{AC}| = \sqrt{(-2-6)^2 + (2+2)^2} = \sqrt{8^2 + 4^2} = \sqrt{64+16} = \sqrt{80} = 4\sqrt{5}$$

$$|\overline{BC}| = \sqrt{(-2-1)^2 + (2+2)^2} = \sqrt{3^2 + 4^2} = \sqrt{9+16} = \sqrt{25} = 5$$

$$\text{Since } |\overline{AB}| = |\overline{BC}| = 5$$

Thus, triangle is an isosceles.

EXAMPLE-2

Express by an equation the fact that the distance from $P(x,y)$ to $A(2,3)$ is twice the distance from $P(x,y)$ to $B(3,4)$

SOLUTION:

Given $A(2,3)$, $B(3,4)$ and $P(x,y)$, where $P(x,y)$ be any point, According to the condition of the question.

$$|AP| = 2 |BP|, \text{ using distance formula.}$$

$$\sqrt{(x-2)^2 + (y-3)^2} = 2\sqrt{(x-3)^2 + (y-4)^2}$$

Taking square on both sides

$$(x-2)^2 + (y-3)^2 = 4[(x-3)^2 + (y-4)^2]$$

$$x^2 - 4x + 4 + y^2 - 6y + 9 = 4[x^2 - 6x + 9 + y^2 - 8y + 16]$$

$$x^2 + y^2 - 4x - 6y + 13 = 4x^2 + 4y^2 - 24x - 32y + 100$$

$$3x^2 + 3y^2 - 20x - 26y + 87 = 0$$

EXAMPLE-3

The vertices of a triangle are $A(1,1)$, $B(5,5)$ and $C(9,1)$. Prove that the triangle is a right triangle.

SOLUTION:

Given $A(1,1)$, $B(5,5)$ and $C(9,1)$. using distance formula,

$$|AB| = \sqrt{(5-1)^2 + (5-1)^2} = \sqrt{4^2 + 4^2} = \sqrt{32}$$

$$|AC| = \sqrt{(9-1)^2 + (1-1)^2} = \sqrt{8^2} = \sqrt{64}$$

$$|BC| = \sqrt{(9-5)^2 + (1-5)^2} = \sqrt{4^2 + 4^2} = \sqrt{32}$$

By Pythagoras theorem,

$$\begin{aligned} |AB|^2 + |BC|^2 &= 32 + 32 \\ &= 64 \\ &= |AC|^2 \end{aligned}$$

Thus ΔABC is a right triangle.

10.2 COLLINEAR POINTS

10.2.1 Collinear Points

Collinear points are points which are the elements of the set of points forming a straight line.

In the given figure (i) the points $A, B, C, D, \dots$ are collinear. If three points are collinear, then one of the points must be lying in between the other two points.




Fig (i)




Fig (ii)

In the figure (ii) 'B' is the point in between the point A and C.

In this case, $|AB| + |BC| = |AC|$.

10.2.2 Collinear and Non-Collinear Points


A line-segment is a subset of a line, consisting of two end points and the set of infinite number of points between them on the line.



In the given figure, $\overline{CD}$ is the line-segment which is a sub-set of a line AB (or $\overline{AB}$). The point C and D are on the line AB and are collinear.

The three or more than three points which are not present on the same straight line are called non-collinear points.

In the given figure P, Q and R are non-collinear points.



Remember that:

In the figure P, Q, R are non-collinear points.

- (i) Two points are always collinear.
- (ii) Three points may or may not be collinear.

10.2.3 Collinearity of Three Points

EXAMPLE-1

Points A, B and C are on the number line at a distance of 1, 4 and 8 units respectively from the origin.

Find $\overline{AB}$, $\overline{BC}$ and $\overline{AC}$, and show that $\overline{AB} + \overline{BC} = \overline{AC}$

SOLUTION:

$$|AB| = 4 - 1$$

$$|AB| = 3$$


$$|BC| = 8 - 4 = 4$$

$$|AC| = 8 - 1 = 7$$

$$\text{Now } |AB| + |BC| = 3 + 4$$

$$= 7 = |AC|$$

$$\text{Thus } |AB| + |BC| = |AC|$$



EXAMPLE-2

Show that the points $A(1,4)$, $B(5,6)$ and, $C(9,8)$ are collinear.

SOLUTION:

Given $A(1,4)$, $B(5,6)$ and, $C(9,8)$.

Using distance formula, we have.

$$|AB| = \sqrt{(5-1)^2 + (6-4)^2} = \sqrt{4^2 + 2^2} = \sqrt{20} = 2\sqrt{5}$$

$$|BC| = \sqrt{(9-5)^2 + (8-6)^2} = \sqrt{4^2 + 2^2} = \sqrt{20} = 2\sqrt{5}$$

$$|AC| = \sqrt{(9-1)^2 + (8-4)^2} = \sqrt{8^2 + 4^2} = \sqrt{80} = 4\sqrt{5}$$

$$\text{Now } |AB| + |BC| = 2\sqrt{5} + 2\sqrt{5}$$

$$= 4\sqrt{5}$$

$$= |AC|$$

Thus, the points A, B , and C are collinear.

EXAMPLE-3

Show that the points $A(4,3)$, $B(-2,3)$ and $C(-6,3)$ are collinear.

SOLUTION:

Given $A(4,3)$, $B(-2,3)$ and $C(-6,3)$.

Using distance formula, we have.

$$|AB| = \sqrt{(-2-4)^2 + (3-3)^2} = \sqrt{36+0} = 6$$

$$|BC| = \sqrt{(-6-2)^2 + (3-3)^2} = \sqrt{16+0} = 4$$

$$|AC| = \sqrt{(-6-4)^2 + (3-3)^2} = \sqrt{100} = 10$$

$$\text{Now } |AB| + |BC| = 6 + 4$$

$$= 10$$





$$= |AC|$$

Thus, the points A, B , and C are collinear.

10.2.4 Use of Distance Formula (for The Non-collinear Points)

We use the distance formula to show that the given three non-collinear points form:-


- ▶ a right angle triangle
- ▶ an isosceles triangle
- ▶ an equilateral triangle
- ▶ a scalene triangle



EXAMPLE-1

Show that the points $A(-1,2)$, $B(7,5)$ and $C(2,-6)$ are vertices of a right triangle.

SOLUTION: Given $A(-1,2)$, $B(7,5)$, $C(2,-6)$.



Let a, b, c denote the lengths of the sides $\overline{BC}$, $\overline{CA}$, and $\overline{AB}$ respectively of ΔABC , using distance formula

$$|PQ| = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

we have

$$a = |\overline{BC}| = \sqrt{(2-7)^2 + (-6-5)^2} = \sqrt{5^2 + 11^2} = \sqrt{146}$$

$$b = |\overline{CA}| = \sqrt{(2-(-1))^2 + (-6-2)^2} = \sqrt{3^2 + 8^2} = \sqrt{73}$$

$$c = |\overline{AB}| = \sqrt{(7-(-1))^2 + (5-2)^2} = \sqrt{(8)^2 + (3)^2} = \sqrt{64+9} = \sqrt{73}$$

$$\text{clearly } |\overline{AB}|^2 + |\overline{CA}|^2 = c^2 + b^2$$

$$= 73 + 73 = 146 = a^2$$

$$= |\overline{BC}|^2$$

Thus, ΔCAB , is a right triangle with right angle at A .

EXAMPLE-2

Show that the points $A(3,1)$, $B(-2,-3)$ and $C(2,2)$ are vertices of an Isosceles triangle.

SOLUTION: Given $A(3,1)$, $B(-2,-3)$ and $C(2,2)$.

Let $\overline{AB} = c$, $\overline{BC} = a$ and $\overline{CA} = b$ be the lengths of the sides of a ΔABC .

Using distance formula, we have

$$a = |\overline{BC}| = \sqrt{(2-(-2))^2 + (2+3)^2} = \sqrt{4^2 + 5^2} = \sqrt{41}$$

$$b = |\overline{CA}| = \sqrt{(3-2)^2 + (1-2)^2} = \sqrt{1^2 + 1^2} = \sqrt{2}$$

$$c = |\overline{AB}| = \sqrt{(-2-3)^2 + (-3-1)^2} = \sqrt{5^2 + 4^2} = \sqrt{41}$$

$$|\overline{AB}| = |\overline{BC}|$$

$$\text{Here } c = a = \sqrt{41}$$

That is, the two sides are equal in length.

Thus, ΔABC is an Isosceles triangle.

EXAMPLE-3

Show that the points $A(-3,0)$, $B(3,0)$ and $C(0,3\sqrt{3})$ are the vertices of an equilateral triangle.

SOLUTION: Given $A(-3,0)$, $B(3,0)$ and $C(0,3\sqrt{3})$.

Using distance formula, we have,

$$|AB| = \sqrt{(-3-3)^2 + (0-0)^2} = \sqrt{(-6)^2} = \sqrt{36} = 6$$

$$|BC| = \sqrt{(3-0)^2 + (0-3\sqrt{3})^2} = \sqrt{9+27} = \sqrt{36} = 6$$

$$|AC| = \sqrt{(-3-0)^2 + (0-3\sqrt{3})^2} = \sqrt{9+27} = \sqrt{36} = 6$$

Here $|AB| = |BC| = |AC| = 6$

That is, three sides of ΔABC are equal in length.

Thus, ΔABC is an equilateral triangle.

EXAMPLE-4

Show that the points $A(5,3)$, $B(-2,2)$ and $C(4,2)$ are vertices of a scalene triangle.

SOLUTION: Given $A(5,3)$, $B(-2,2)$ and $C(4,2)$.

Let $\overline{BC} = a$, $\overline{CA} = b$, $\overline{AB} = c$ be the lengths of the sides of a ΔABC .

Using distance formula, we have

$$|BC| = a = \sqrt{(4+2)^2 + (2-2)^2} = \sqrt{6^2} = 6$$

$$|CA| = b = \sqrt{(5-4)^2 + (3-2)^2} = \sqrt{1^2 + 1^2} = \sqrt{2}$$

$$|AB| = c = \sqrt{(-2-5)^2 + (2-3)^2} = \sqrt{7^2 + 1^2} = \sqrt{50} = 5\sqrt{2}$$

As, $|AB| = c$, $|BC| = a$, $|CA| = b$ are all different in length.

Thus ΔABC is a scalene triangle.

EXERCISE - 10.1

- 1- Describe the location of these points on the number plane.
(i) $(1,0)$ (ii) $(0,4)$ (iii) $(-2,4)$ (iv) $(3,6)$
(v) $(-4,0)$ (vi) $(-8,-8)$ (vii) $(7,-5)$ (viii) $(-8,10)$
(ix) $(0,-7)$ (x) $(8,-3)$
- 2- Find the distance between the following pairs of points.
(i) $(2,1), (-4,3)$ (ii) $(-1,3), (-2,-1)$
(iii) $(7,-2), (-2,3)$ (iv) $(a,-b), (b,-a)$
- 3- Express by an equation, the fact, that the point $P(x,y)$ is equidistant from $A(2,4)$ and $B(6,8)$.
- 4- Show that the points $A(5,4)$, $B(4,-3)$, $C(-2,5)$ are equidistant from $D(1,1)$.
- 5- Find the point on the x -axis which is equidistant from $(2,4)$ and $(6,8)$.
(Hint: call the point $(x,0)$. Find x .)
- 6- Show that the points $A(0,2)$, $B(3,-2)$ and $C(0,-2)$ are vertices of a right triangle.
- 7- Show that the points $A(-1,1)$, $B(3,2)$, $C(7,3)$ are collinear.
- 8- Show that the points $A(6,1)$, $B(2,7)$ and $C(-6,-7)$ are vertices of a right triangle.
- 9- Show that the points $A(2,4)$, $B(6,2)$, $C(4,3)$ are collinear.
- 10- Show that the points $A(4,-2)$, $B(-2,4)$ and $C(5,5)$ are vertices of an isosceles triangle.
- 11- Show that the points $A(-2,11)$, $B(-6,-3)$ and $C(4,-9)$ are of a scalene triangle.
- 12- Show that the points $A(6,1)$, $B(2,7)$ and $C(-6,7)$ are of a scalene triangle.
- 13- Show that the points $A(2,-5)$, $B(-4,-3)$ and $C(-1,5)$ are of an equilateral triangle.

Review Exercise-10

I- Encircle the Correct Answer.

- $d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$ is called
 - distance formula
 - collinear points
 - non-collinear points
 - equal points
- A point in a cartesian plane determines a unique ordered pair of:
 - set
 - abscissa
 - numbers
 - ordinate
- In the plane with every ordered pair is associated:
 - a unique point
 - zero
 - two points
 - four points
- Points lying on the same line are called:
 - non-collinear
 - collinear
 - equal
 - overlapping
- Points which do not lie on the same straight line are called:
 - non-collinear
 - collinear
 - equal
 - zero
- Point on the axis do not lie in any:
 - a plane
 - line
 - quadrant
 - circle
- The co-ordinates of the origin are:
 - 0
 - (1,0)
 - (0,0)
 - (0,1)
- Points on the negative x-axis have negative:
 - abscissa
 - ordinate
 - value
 - fraction
- A point in 4th quadrant has its ordinate:
 - positive
 - negative
 - zero
 - one
- A point in the first quadrant is characterized by the fact that both its co-ordinates are:
 - zero
 - positive
 - negative
 - positive and negative oth

II- Fill in the blanks.

1. $d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$ is called _____.
2. A point in a cartesian plane determines a _____ ordered pair of numbers.
3. With every ordered pair is associated a _____ point in the plane .
4. Points lying on the same line are called _____ points.
5. Points which do not lie on the same straight line are called _____ points.
6. Points on the axes do not lie in any _____.
7. The origin has the co-ordinates _____.
8. Points on the negative x-axis have negative abscissa and their ordinate is _____.
9. A point in the 4th quadrant has its abscissa positive and its ordinate _____.
10. A point in the first quadrant is characterized by the fact, that both its co-ordinates are _____.

SUMMARY

Distance Formula: $d = |\overline{PQ}| = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$

- 1- A point in a number plane determines a unique ordered pair of numbers.
- 2- With every ordered pair of numbers a unique point is associated in the plane.

Collinear points: Points lying on the same straight line are called collinear points.

Non-Collinear points: Points which do not lie on a same straight line are called non-collinear points.

ANSWERS

Exercise 1.1

- 1- 9,8 2- -11 3- 9 4- 11,71 5- 1,0 6- 18.86 7- 804.57 8- 25,1
- 9- $\frac{2y}{3x^2}$ 10- $\frac{25a}{14b^2}$ 11- $\frac{4a^3b^3}{5a^2+3b}$ 12- $\frac{2m}{3x^5-4m^2x^3}$ 13- $\frac{5}{c+d}$
- 14- $\frac{x+y}{-3}$ 15- $\frac{2x^3-x^2y+xy^2}{x^3-x^2y+xy^2-y^3}$ 16- $\frac{4x^2-2x}{x^2-1}$ 17- $\frac{3x-1}{x^3-7x-6}$
- 18- $\frac{2x-3y}{2x+3y}$ 19- $\frac{x-2y}{x^2-y^2}$ 20- $\frac{x-2y}{xy-y^2}$ 21- $\frac{2}{x-1}$
- 22- $\frac{37x+1}{x^2-12x+27}$ 23- $\frac{x^2-4x+4}{x^2+2x}$ 24- $\frac{-(x+6)}{x+1}$ 25- $\frac{x^3+x^2+20x}{x^2+4x-5}$
- 26- $\frac{3x+4}{2x+1}$ 27- $\frac{4x^3-x}{2x^2-1}$ 28- $\frac{x}{x^3-2x^2+2x-1}$ 29- $\frac{x}{3x-9}$
- 30- x 31- 1 32- $x-1$

Exercise 1.2

- 1- $2x^2+8y^2$ 2- $50x^2+18y^2$ 3- $24lm$ 4- ℓ^8-m^8 5- $a^3b^3-\frac{1}{a^3b^3}-3ab+\frac{3}{ab}$
- 6- $4x^2+9y^2+4+12xy+12y+8x$ 7- $8p^3+12p^2q+6pq^2+q^3$
- 8- $9p^2+q^2+r^2+6pq+2qr+6pr$ 9- $8x^3+36x^2y+54xy^2+27y^3$
- 10- $(x+y-1)(x^2+y^2+2xy+x+y+1)$ 11- $(x-y+4)(x^2+y^2-2xy-4x+4y+16)$
- 12- $(2x+3y)(4x^2-6xy+9y^2)$ 13- $(x+3y)(x^2-3xy+9y^2)(x-3y)(x^2+3xy+9y^2)$
- 14- $(2a+b)(2a-b)(4a^2-2ab+b^2)(4a^2+2ab+b^2)$ 15- 4 17- 17,4 18- 14
- 19- 133 20- 118 21- 20 22- 46

Exercise 1.3

- 1- (i) $\frac{\sqrt{5}}{5}$, (ii) $\frac{7\sqrt{6}}{3}$, (iii) $\frac{\sqrt{42}}{7}$ 2- (i) $3\sqrt{2}$, (ii) $35\sqrt{2}$, (iii) $4\sqrt{15} - 6\sqrt{6} - 2\sqrt{10} + 6$
 (iv) $30 - 6\sqrt{5} + 5\sqrt{2} - \sqrt{10}$ (v) $5\sqrt{3} - \sqrt{15} - 10 + 2\sqrt{5}$ (vi) $35 + 7\sqrt{2} + 5\sqrt{3} + \sqrt{6}$
 3- (i) $2 - \sqrt{3}$ (ii) $\frac{4 + \sqrt{5}}{11}$ (iii) $2\sqrt{3}(\sqrt{7} - \sqrt{5})$ (iv) $\frac{x + y - 2\sqrt{xy}}{x - y}$
 (v) $\frac{105 - 10\sqrt{7}}{59}$ (vi) $5 + 2\sqrt{6}$ (vii) $\frac{29(11 - 3\sqrt{5})}{76}$ (viii) $\frac{3\sqrt{7} - 2\sqrt{3}}{3}$
 4- (i) $2\sqrt{5}$ (ii) 18 5- (i) $2\sqrt{3}$ (ii) 14 6- (i) $-2\sqrt{2}$ (ii) 10
 7- (i) $\frac{24 - 6\sqrt{2}}{7}$ (ii) $\left(\frac{-18 + 8\sqrt{2}}{7}\right)$ 8- (i) 40 (ii) 36
 9- (i) $\frac{2b^2 - a^2 + 2b\sqrt{b^2 - a^2}}{a^2}$ (ii) $\frac{a - \sqrt{a^2 - 9}}{3}$

Review Exercise 1

I- Encircle the Correct Answer.

- 1- b 2- b 3- d 4- c 5- d 6- a 7- d 8- b 9- c 10- a

II- Fill in the blanks.

- 1- rational number 2- rational expression 3- $4ab$ 4- $2(a^2 + b^2)$
 5- $(a+b)^3$ 6- $(a-b)^3$ 7- $a^3 - b^3$ 8- $a^3 + b^3$ 9- surd 10- 2

Exercise 2.1

- 1- $(x+y)(3a-7b)$ 2- $(a-x)(x+y)$ 3- $(a-3)(a^2+1)$
 4- $(x-1)(x^2+x-y)$ 5- $(x+2y)(3a-4b)$ 6- $(a-b)(2a+c)$
 7- $(a-b)(a+c)$ 8- $(4-a^3)(2-a)$ 9- $(4x-3a)^2$
 10- $(1-7x)^2$ 11- $5(2x-1)^2$ 12- $2ab(a-b)^2$
 13- $\left(x + \frac{1}{2}\right)^2$ 14- $\left(x - \frac{1}{x}\right)^2$ 15- $5x(x-3)^2$ 16- $(a+b)(a+b+2c)$

ANSWERS

Exercise 2.2

1- $(x+y+a)(x+y-a)$

2- $(2a+b+3c)(2a+b-3c)$

3- $(x+3a+4b)(x+3a-4b)$

4- $(y+x-c)(y-x+c)$

5- $(x+y+2xy)(x+y-2xy)$

6- $(a-2b-3ac)(a-2b+3ac)$

7- $(x-y-a+b)(x-y+a-b)$

8- $(y^2+2y+2)(y^2-2y+2)$

9- $(z^2+8y^2-4yz)(z^2+8y^2+4yz)$

10- $(x^3-6x+18)(x^3+6x+18)$

11- $(z^2-3z+4)(z^2+3z+4)$

12- $(2x-y)(x-y)(2x+y)(x+y)$

Exercise 2.3

1- $(x+4)(x+5)$

2- $(x-2)(x+7)$

3- $(x-1)(x+6)$

4- $(x-3)(x-4)$

5- $(x-13)(x+12)$

6- $(x-2)(x+1)$

7- $(x-15)(x+6)$

8- $(a-17)(a+5)$

9- $(7-x)(x+14)$

10- $(y-19)(y+8)$

11- $(x+1)(2x+1)$

12- $(x+1)(3x+2)$

13- $(x-1)(2x+1)$

14- $(2x+3)(3x-1)$

15- $(x+2)(1-2x)$

16- $(2-x)(4+5x)$

17- $(u-2)(3u-4)$

18- $(2x-3)(5x+4)$

19- $(x-6)(5x-2)$

20- $(4x-\sqrt{3})(\sqrt{3}x+2)$

Exercise 2.4

1- $(2x-y)(4x^2+2xy+y^2)$

2- $(3x+1)(9x^2-3x+1)$

3- $(1-7x)(1+7x+49x^2)$

4- $(ab+8)(a^2b^2-8ab+64)$

5- $(3-10y)(9+30y+100y^2)$

6- $(3x-4y)(9x^2+12xy+16y^2)$

7- $(xy+z)(x^2y^2-xyz+z^2)$

8- $(6p-7)(36p^2+42p+49)$

9- $(2x-\frac{1}{3})(4x^2+\frac{2}{3}x+\frac{1}{9})$

10- $(a+b)[a^2-ab+b^2+1]$

11- $(a-b)[1-(a^2+ab+b^2)]$

12- $x(1-2y)(1+2y+4y^2)$

13- $(x-y)(x+y)(x^2+y^2)(x^2+xy+y^2)(x^2-xy+y^2)(x^4-x^2y^2+y^4)$

ANSWERS

$$14- \left(1 - \frac{4p}{q}\right) \left(1 + \frac{4p}{q} + \frac{16p^2}{q^2}\right)$$

$$15- (1+4u)(1-4u+16u^2)$$

$$16- (2x+3y)(4x^2+9y^2-6xy-3)$$

$$17- (z+5)(z^2-5z+25)$$

$$18- (x+y)(x^2-xy+y^2)(x^6-x^3y^3+y^6)$$

$$19- (m+n)(m-n)(m^2+mn+n^2)(m^2-mn+n^2)$$

$$20- x(2x-a)(2x+a)(4x^2+2ax+a^2)(4x^2-2ax+a^2)$$

$$21- (x-3a)(x^2+3ax+9a^2)$$

$$22- (x+3a)(x^2-3ax+9a^2)$$

Exercise 2.5

$$1- 3 \quad 2- -6 \quad 3- -47 \quad 4- 0 \quad 5- -84 \quad 6- \text{yes} \quad 7- \text{yes} \quad 8- \text{no} \quad 9- \text{no}$$

$$10- \text{yes} \quad 11- \text{no} \quad 12- \text{yes} \quad 13- \text{yes} \quad 14- \text{no} \quad 15- \text{no} \quad 16- \text{yes} \quad 17- \text{yes}$$

$$18- \text{no} \quad 19- k=1 \quad 20- k=1$$

Review Exercise 2

I- Encircle the Correct Answer.

$$1- b \quad 2- c \quad 3- d \quad 4- a \quad 5- c \quad 6- b \quad 7- a \quad 8- a \quad 9- a \quad 10- a$$

II- Fill in the blanks.

$$1- \text{one} \quad 2- \text{two} \quad 3- \text{three} \quad 4- (x-3)(x+3) \quad 5- (x+1)(x+3)$$

$$6- (x+2)(x^2-2x+4) \quad 7- (x-2)(x^2+2x+4) \quad 8- 3 \quad 9- 11 \quad 10- 0$$

Exercise 3.1

$$1- ab$$

$$2- 3qr$$

$$3- 4xy^2z^2$$

$$4- 7ab$$

$$5- 3x^2y^2$$

$$6- 2abc$$

$$7- x+4$$

$$8- x^2-y^2$$

$$9- t+3$$

$$10- x-2$$

$$11- 1+x$$

$$12- x-2$$

$$13- x+1$$

$$14- x(x+3)$$

$$15- 5abc$$

ANSWERS

Exercise 3.2

- 1- $x^2 - x + 1$ 2- $2x^2 + 3x - 2$ 3- $2(x-1)$ 4- $9x(x+3)$ 5- $(x-1)^2(x+1)$ 6- $(x-2)$
7- $(x-1)$ 8- $(3x-5)$ 9- $2x+1$ 10- $(x+3)$

Exercise 3.3

- 1- $420a^4x^4y^4$ 2- $15a^4b^3c^5$ 3- $12abc$ 4- $x^2y^2z^2$
5- $p^2q^2(p-q)(p+q)(p^2+pq+q^2)$ 6- $(x+4)(x-4)(x^2-4x+16)$
7- $(x-2)(x+3)(x+1)(x-1)$ 8- $(y+3)(y-2)(y+3)(y-3)$
9- $(1+y)(1-y)(1-2y)(y^2-y+1)$ 10- $(x-y)(x+y)(x^2+y^2)(x^4+x^2y^2+y^4)$
11- $(x+1)(x^2-x+1)(x^2+x+1)^2$
12- $(x+y)(x^2+y^2)(x-y)(x^2-xy+y^2)(x^4-x^2y^2+y^4)$ 13- $(2x+3)(x+1)^2(x+3)$
14- $x^2(x+3)(x-2)(x-3)$ 15- $(x+y)^2(x+2y)^2$

Exercise 3.4

- 1- x^2+1 ; x^4-1 2- $(x^2-4)(x-3)(x^3-x^2-4x+4)$ 3- $2x^2+1$; $2x^4-x^2-1$
4- $2x^2+3x-2$; $(3x-1)(8x^4+6x^3-15x^2+9x-2)$
5- $(3x^2+8x-3); (2x^2-3x+1)(3x^4+17x^3+27x^2+7x-6)$
6- $(x^2+2x-3); (2x^2-x-5)(2x^4+x^3-20x^2-7x+24)$
7- $(x^3-1); (x-1)(x^4+x^3-x-1)$
9- $x^2-12x+35$ 10- $(6x^2+x-2)$ 11- $x+4$ 12- $(x+1); (x^3+1)(x^4+x^3-x-1)$
14- $x^3-7x^2+16x-12$ 15- $x^4+8x^3+11x^2-32x-60$ 16- x^5-x^4-4x+4

Exercise 3.5

- 1- $\frac{2(2a+1)}{a(a+1)(a+2)}$ 2- $\frac{2ax+x-3a-6a^2}{(x-2a)(x-3a)}$ 3- $2+a^4$ 4- $\frac{1}{x^4+x^2+1}$

ANSWERS

$$5- \frac{2b^2(a-c)}{(a+b)(b+c)}$$

$$6- \frac{6x^3}{x^6-1}$$

$$7- \frac{2a^3}{a^2-b^2}$$

$$8- 1$$

$$9- 1$$

$$10- 1$$

$$11- \frac{a}{a-b}$$

$$12- \frac{a+1}{a+2}$$

Exercise 3.6

$$1- \pm(4x+3y) \quad 2- \pm(x-3)(x-4)(x-5) \quad 3- \pm(x+1)(x+7)(2x-3) \quad 4- \pm(x^2+6x+4)$$

$$5- \pm(4x^2+16x+11) \quad 6- \pm\left(x+\frac{1}{x}-5\right) \quad 7- \pm\left(t+\frac{1}{t}-2\right) \quad 8- \pm\left(x^2+\frac{1}{x^2}-2\right)$$

$$9- \pm(2x^2+3x+4) \quad 10- \pm\left(\frac{3x}{2y}-\frac{1}{2}-\frac{2y}{3x}\right) \quad 11- x=8 \quad 12- t=4, m=10$$

Review Exercise 3

I- Encircle the Correct Answer.

$$1- a \quad 2- c \quad 3- c \quad 4- a \quad 5- b \quad 6- a \quad 7- c \quad 8- c \quad 9- a \quad 10- a$$

II- Fill in the blanks.

$$1- two \quad 2- two \quad 3- H.C.F \quad 4- L.C.M \quad 5- H.C.F \quad 6- second expression$$

$$7- 2x+1 \quad 8- x+2 \quad 9- 2x^2y^3 \quad 10- 6x^2y^2z$$

Exercise 4.1

$$1- (i) 8, (ii) 80, (iii) 11, (iv) 2 \quad 2- \frac{5}{2} \quad 3- 2 \quad 4- -7 \quad 5- -2$$

$$6- 3 \quad 7- 4 \quad 8- 4 \quad 9- 3 \quad 10- \{4\} \quad 11- \{9\} \quad 12- \{18\} \quad 13- \{8\}$$

$$14- \{\} \quad 15- \{\} \quad 16- \{13, 5\} \quad 17- \{8\} \quad 18- \{13\} \quad 19- \{101\} \quad 20- \{15\}$$

Exercise 4.2

$$1- \pm 9 \quad 2- -1, 7 \quad 3- -6, 4 \quad 4- -1, 4 \quad 5- \frac{5}{3}, \frac{-13}{3} \quad 6- x < 7$$

ANSWERS

- 7- $x > -3$ 8- $x < -1$ 9- $x < -10$ 10- $x > -\frac{17}{9}$ 11- $x < -21$
12- $x > -12\frac{5}{7}$ 13- $x \geq 6$ 14- $x \leq 1\frac{7}{18}$ 15- $x \geq 1\frac{1}{2}$ 16- $x \geq 0$

Review Exercise 4

I- Encircle the Correct Answer.

- 1- a 2- c 3- c 4- c 5- c 6- a 7- c 8- a

II- Fill in the Blanks.

- 1- $>$ 2- $>$ 3- $<$ 4- $<$ 5- $>$ 6- $>$ 7- $>$ 8- $<$
9- $<$ 10- $<$ 11- $<$ 12- $>$

Exercise 5.1

- 1- -2,6 2- 1,5 3- -8,1 4- 2,3 5- $2\frac{4}{3}$ 6- $-8\frac{1}{2}$ 7- 3,-4
8- $3, -\frac{1}{3}$ 9- $2, -\frac{1}{2}$ 10- $2, -\frac{4}{5}$ 11- $2, -\frac{3}{2}$ 12- $-\frac{1}{2}, \frac{4}{5}$ 13- $1, \frac{-1}{2}$
14- $5 \pm 2\sqrt{7}$ 15- $3 \pm 2\sqrt{3}$ 16- $\frac{-1 \pm \sqrt{5}}{2}$ 17- $-3 \pm 2\sqrt{3}$ 18- $\frac{2 \pm \sqrt{2}}{2}$
19- $\frac{3 \pm \sqrt{3}}{2}$ 20- $\frac{-5 \pm \sqrt{73}}{6}$ 21- $\frac{-m \pm \sqrt{m^2 - 4n}}{2}$ 22- $\frac{3 \pm 4\sqrt{15}}{11}$ 23- $-2 \pm \sqrt{17}$
24- $\frac{10 \pm 4\sqrt{15}}{5}$ 25- $\{13, -2\}$

Exercise 5.2

- 1- 2,3 2- $\frac{3}{4}, \frac{1}{2}$ 3- $-1, \frac{2}{3}$ 4- $-1, \frac{3}{2}$ 5- -5,3 6- $3, -\frac{7}{2}$ 7- $\pm\sqrt{10}$
8- $\pm 2\sqrt{6}$ 9- ± 8 10- $\frac{5}{3}$ 11- 0,-5 12- $5, \frac{1}{3}$

ANSWERS

Exercise 5.3

- 1- 5, 7 2- 8, 10 3- 9, 18 4- 5 5- 5, 6 6- 12, 13 7- 7, 9
8- 4, 8 or 8, 4

Review Exercise 5

I- Encircle the Correct Answer.

- 1- a 2- b 3- b 4- a 5- c 6- c 7- c 8- c 9- b 10- b

II- Fill in the blanks.

- 1- quadratic 2- quadratic formula 3- $x(2x - 3)$ 4- $\{-1, 3\}$ 5- three
6- quadratic formula 7- $\{2, 3\}$ 8- $\{\pm 3\}$ 9- $(x-2)(x+2)(x^2+4)$ 10- $\{\pm 1\}$

Exercise 6.1

- 1- $2-by-2, 3-by-1, 3-by-2$ 2- $2-by-2, 3-by-3, 1-by-3$
3- 5 4- $B=F, G=J, H=K, C=E, A=D$

Exercise 6.2

- 1- Row matrix = A, Column matrix = C, Square matrices = B, D, E, F Rectangular matrices = A, C, G
2- Diagonal matrix are A, B, C, D, E, F, G Scalar matrix are B, D, E, G, Identity is D

$$3- \begin{bmatrix} 3 & -1 \\ 4 & 4 \end{bmatrix}, \begin{bmatrix} -3 & -1 \\ -2 & 4 \end{bmatrix}, \begin{bmatrix} a & c \\ -b & d \end{bmatrix}, \begin{bmatrix} l & p & a \\ m & q & b \\ n & r & c \end{bmatrix} \quad 4- A, C \quad 5- A, C, E \quad 6- C \quad 7- A$$

Exercise 6.3

$$1- (i) \begin{bmatrix} 2 & 4 & 9 \\ 3 & 8 & 11 \\ 5 & 13 & 1 \end{bmatrix} \quad (ii) \begin{bmatrix} 2 & 2 & -1 \\ -1 & 2 & -1 \\ 3 & 5 & 5 \end{bmatrix} \quad (iii) \begin{bmatrix} -2 & -2 & 1 \\ 1 & -2 & 1 \\ -3 & -5 & -5 \end{bmatrix} \quad (iv) \begin{bmatrix} 4 & 9 & 23 \\ 8 & 19 & 28 \\ 11 & 30 & 0 \end{bmatrix}$$

$$(v) \begin{bmatrix} 6 & 5 & -8 \\ -5 & 3 & -9 \\ 8 & 11 & 17 \end{bmatrix} \quad (vi) \begin{bmatrix} 2 & 1 & -6 \\ -3 & -1 & -7 \\ 2 & 1 & 7 \end{bmatrix} \quad 2- -A = \begin{bmatrix} -4 & -3 \\ -2 & -6 \end{bmatrix}, -B = \begin{bmatrix} -\sqrt{2} & -3 \\ -4 & -\sqrt{3} \end{bmatrix}$$

ANSWERS

$$-C = \begin{bmatrix} -1 \\ 7 \\ -4 \end{bmatrix}, -D = \begin{bmatrix} -1 & 0 & 2 \\ 0 & -3 & -4 \\ -2 & 1 & 3 \end{bmatrix}, -E = [-2 \ -5 \ 3] \quad 4. -1, 2 \quad 6. X = \begin{bmatrix} 2 & 1 \\ -4 & 2 \\ -3 & 2 \end{bmatrix}$$

$$7. a=2, b=-4, c=4, d=3, e=4, f=2 \quad 8. w=-1, x=1, y=7, z=-8$$

$$9. \begin{bmatrix} -a & -b \\ -c & -d \end{bmatrix}$$

Exercise 6.4

$$9. [12 \ 13]$$

$$10. \begin{bmatrix} 5 \\ -3 \end{bmatrix}$$

$$11. \begin{bmatrix} 2 & 4 \\ 1 & -5 \end{bmatrix}$$

$$12. \begin{bmatrix} 1 & -9 \\ -3 & 17 \end{bmatrix}$$

$$13. \begin{bmatrix} 10 & 1 \\ -2 & 10 \end{bmatrix}$$

$$14. \begin{bmatrix} 10 & -14 \\ 15 & 3 \end{bmatrix}$$

$$15. a = \frac{10}{7}, b = 0$$

Exercise 6.5

$$1. (i) uy - vx \quad (ii) -13 \quad (iii) 0 \quad (iv) \frac{13}{64}$$

$$2. (i) \text{ singular} \quad (ii) \text{ non-singular} \quad (iii) \text{ non-singular}$$

$$3. (i) \begin{bmatrix} 3 & -2 \\ -1 & 1 \end{bmatrix} \quad (ii) \begin{bmatrix} 3 & -1 \\ -5 & 2 \end{bmatrix} \quad (iii) \begin{bmatrix} \frac{1}{2} & 0 \\ \frac{1}{6} & \frac{1}{3} \end{bmatrix}$$

(iv) Inverse does not exist

$$(v) \begin{bmatrix} 4 & \frac{-3}{2} \\ -1 & \frac{1}{2} \end{bmatrix}$$

$$(vi) \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix}$$

$$(vii) \begin{bmatrix} \frac{3}{5} & \frac{4}{5} \\ \frac{-4}{5} & \frac{3}{5} \end{bmatrix}$$

$$4. M^{-1} = \begin{bmatrix} -2 & 1 \\ 3 & -1 \\ 2 & 2 \end{bmatrix}$$

Exercise 6.6

$$1. \frac{-9}{2} \quad 2. (i) (3, 1) \quad (ii) \left(\frac{7}{3}, \frac{3}{2}\right) \quad (iii) \left(\frac{-1}{2}, \frac{2}{5}\right) \quad (iv) (-1, 5) \quad (v) (0, 2) \quad (vi) (-2, 6)$$

$$3. (3, -1) \quad 4. (i) (-1, 2) \quad (ii) (1, -1) \quad (iii) (4, -1) \quad (iv) \text{No Solution} \quad (v) (2, -1)$$

$$(vi) \left(\frac{31}{21}, \frac{59}{21}\right)$$

$$5. (i) 2x - y = 2, 5x + 2y = 4$$

$$(ii) -5x + 2y = 2, 2x - 3y = -1$$

$$(iii) -4x + y = 1, 5x + 4y = -1$$

$$(iv) 0.8x - 0.6y = 1, 0.6x + 0.8y = 2$$

ANSWERS

Review Exercise 6

I- Encircle the Correct Answer.

- 1- a 2- a 3- a 4- c 5- a 6- c 7- b 8- c 9- c 10- c

II- Fill in the blanks.

- 1- order 2- row matrix 3- same order 4- same 5- equal 6- 1
7- associative 8- skew symmetric 9- $B^t A^t$ 10- $B^{-1} A^{-1}$

Exercise 7.1

- 1- (i) 130° (ii) 115° (iii) 42° (iv) 30° (v) 108° (vi) 20° 2- $105^\circ, 75^\circ$ 3- 70°
4- $-0^\circ, 100^\circ$ 5- $70^\circ, 30^\circ$ 6- $x + 90^\circ + 30^\circ = 180^\circ \Rightarrow x = 60^\circ$ 7- (i) $a = 40^\circ$
(ii) $c = 35^\circ, d = 145^\circ$ (iii) $e = 29^\circ, f = 151^\circ$ (iv) $b = 135^\circ$
(v) $q = 77^\circ, P = 103^\circ, r = 103^\circ$ (vi) $j = 30^\circ, k = 150^\circ, l = 30^\circ$
(vii) $g = 140^\circ, h = 40^\circ, i = 140^\circ$ (viii) $k = 145^\circ$
(ix) $P = 58^\circ, M = 122^\circ, N = 122^\circ$ (x) $a = 158^\circ, b = 112^\circ$

Exercise 7.2

- 1- (a) $(\angle 1, \angle 2), (\angle 3, \angle 4)$ (b) $(\angle 1, \angle 6), (\angle 3, \angle 8), (\angle 2, \angle 7), (\angle 5, \angle 4)$ (c) none
(d) $(\angle 1, \angle 8), (\angle 1, \angle 4), (\angle 4, \angle 7), (\angle 7, \angle 8), (\angle 5, \angle 6), (\angle 5, \angle 2), (\angle 2, \angle 3), (\angle 3, \angle 6)$
(e) $(\angle 1, \angle 7), (\angle 4, \angle 8), (\angle 5, \angle 3), (\angle 2, \angle 6)$
2- (a) $(\angle 1, \angle n), (\angle m, \angle r)$ (b) $(\angle p, \angle n), (\angle m, \angle s), (\angle q, \angle r), (\angle 1, \angle t)$ (c) none
(d) $(\angle p, \angle m), (\angle n, \angle s), (\angle q, \angle t), (\angle r, \angle t), (\angle q, \angle p), (\angle 1, \angle m), (\angle r, \angle n), (\angle t, \angle s)$
(e) $(\angle p, \angle t), (\angle m, \angle q), (\angle n, \angle t), (\angle s, \angle r)$

Exercise 7.3

- 1- yes, no, yes 2- yes 3- yes 4- 10cm, 12cm, 14cm, 16cm, 18cm
5- 6cm, 12cm, 18cm, 21cm 6- 15cm, 21cm, 9cm, 12cm, 1:3
7- $\overline{AB} \cong \overline{DE}, \overline{AC} \cong \overline{DF}, \overline{BC} \cong \overline{EF}$ $\angle A \cong \angle D, \angle B \cong \angle E, \angle C \cong \angle F$
8- No: size may be different 9- yes: size and shape are same

Exercise 7.4

- 1- (a) (i) $\overline{AB} \cong \overline{FD}$ (ii) $\overline{BC} \cong \overline{DE}$ (iii) $\overline{AC} \cong \overline{FE}$ (iv) $\angle A \cong \angle F$ (v) $\angle B \cong \angle D$ (vi) $\angle C \cong \angle E$
(b) $\angle R$ (c) $\overline{EF}$ (d) $S.A.S \cong S.A.S$ (e) $ASA \cong ASA$
2- (i) $\triangle ABC \cong \triangle DFE$ by $S.S.S \cong S.S.S$ (ii) $\triangle XYZ \cong \triangle DFE$ by $S.S.A \cong S.S.A$
(iii) $\triangle ABC \cong \triangle CDA$, by $A.S.A \cong A.S.A$ (iv) $\triangle PQT \cong \triangle SRT$ by $S.A.S \cong S.A.S$

ANSWERS

- 3- $\overline{AD} \cong \overline{DA}$, $\overline{DB} \cong \overline{AC}$, $\overline{AB} \cong \overline{DC}$, $\angle BAD \cong \angle CDA$, $\angle ADB \cong \angle DAC$, $\angle ABD \cong \angle DCA$,
Condition used $S.S.S \cong S.S.S$, $m\angle ADB = 40^\circ$
- 4- (i) Similar Triangles (ii) Similar Parallelogram (iii) Similar Triangles
- 5- $\overline{MN} \Leftrightarrow \overline{PQ}$, $\overline{NO} \Leftrightarrow \overline{QR}$, $\overline{PR} \Leftrightarrow \overline{MO}$, $\angle 1 \Leftrightarrow \angle 4$

Exercise 7.5

- (i) rectangle (ii) square (iii) quadrilateral (iv) bisect (v) congruent

Exercise 7.6

- (i) circle (ii) radius (iii) chord (iv) diameter (v) semicircle
(vi) major arc (vii) radius (viii) sector (ix) secant line (x) right angle

Review Exercise 7

I- Encircle the Correct Answer.

- 1- b 2- c 3- b 4- b 5- a 6- c 7- a 8- b 9- c 10- c

II- Fill in the blanks.

- 1- adjacent 2- supplementary 3- obtuse 4- vertical 5- 180° 6- each other
7- congruent 8- scalene 9- diameter 10- right

Review Exercise 8

I- Encircle the Correct Answer.

- 1- c 2- c 3- c 4- c 5- a 6- a 7- a 8- c 9- c 10- a

II- Fill in the blanks.

- 1- concurrent 2- concurrent 3- concurrent 4- concurrent 5- altitude
6- median 7- angle bisector 8- three 9- three 10- three

Exercise 9.1

- 1- (i) 5 (ii) 12 (iii) $4\sqrt{231}$ 3- $\sqrt{2}l$ 4- $8\sqrt{2}$
5- (i) right triangle (ii) not right Δ (iii) right Δ 6- 15cm 7- 7cm 8- 8m 9- 5

ANSWERS

Exercise 9.2

- 1- 20 stones 2- 24000 stones 3- Rs. 223 4- 645.50m 5- 1mm 54sec
 6- 98 cm^2 7- (i) 8967 cm^2 (ii) 16.8 m^2 8- 9000 m^2 9- $16\sqrt{110}\text{ m}^2$
 10- (i) 44 cm^2 (ii) 0.5 (iii) 401.14 mm^2 11- 154 m^2 12- $16\sqrt{3}\text{ m}^2$ 13- $9\sqrt{3}\text{ cm}^2$
 14- 210 cm^2 15- 7cm, 21cm 16- 1666.67cm 17- 72cm 18- 3600 cm^2 19- 4cm

Exercise 9.3

- 1- 64 cm^3 2- 64 cm^3 3- 24 m^3 4- 502.86 cm^3 5- 94.3 cm^3 6- 113.1 cm^3
 7- 127.3 cm^3 8- 339.4 cm^3

Review Exercise 9

I- Encircle the Correct Answer.

- 1- a 2- c 3- c 4- a 5- d 6- c 7- a 8- c 9- c

II- Fill in the blanks.

- 1- Pythagoras 2- Area 3- $\frac{1}{2} \times \text{base} \times \text{altitude}$ 4- $\sqrt{S(s-a)(s-b)(s-c)}$
 5- $\frac{\sqrt{3}a^2}{4}$ 6- $l \times b$ 7- πr^2 8- l^3 9- $l \times b \times h$ 10- $\frac{1}{3} \pi r^2 h$

Exercise 10.1

- 1- (i) lies on $\overrightarrow{OX}$ (ii) lies on $\overrightarrow{OY}$ (iii) lies in QII (iv) in QI
 (v) on $\overrightarrow{OX}$ (vi) in $QIII$ (vii) in QIV (viii) in QII
 (ix) on $\overrightarrow{OY}$ (x) in QIV
 2- (i) $2\sqrt{10}$ (ii) $\sqrt{17}$ (iii) $\sqrt{106}$ (iv) $(a-b)\sqrt{2}$ 3- $x + y - 10 = 0$
 5- (10,0)

Review Exercise 10

I- Encircle the Correct Answer.

- 1- a 2- c 3- a 4- b 5- a 6- c 7- c 8- a 9- b 10- b

II- Fill in the blanks.

- 1- Distance formula 2- unique 3- unique 4- collinear 5- non-collinear
 6- quadrant 7- (0,0) 8- zero 9- negative 10- positive

GLOSSARY

Unit-1 ALGEBRAIC FORMULAS AND APPLICATIONS

Formula: Where we have a rule to calculate some quality, we write the rule as a formula.

$$(a \pm b)^2 = a^2 \pm 2ab + b^2$$

$$(a+b)^2 + (a-b)^2 = 2(a^2 + b^2)$$

$$(a+b)^2 - (a-b)^2 = 4ab$$

$$(a+b+c)^2 = (a^2 + b^2 + c^2 + 2ab + 2bc + 2ac)$$

$$(a \pm b)^3 = a^3 \pm 3ab(a \pm b) \pm b^3$$

$$(x+y)(x-y)(x^2 - xy + y^2)(x^2 - xy + y^2)$$

Surd: A surd is an irrational number that contains an irrational square root.

Pure Surd: A surd which has unity only as rational factor, the other factor being irrational is called a pure surd.

Mixed surd: A surd which has rational factor other than unity, the other factor being irrational is called mixed surd.

Similar surd: Surds having the same irrational factor are called similar or like surd.

Unlike surd: Surd having no common irrational factor are known as unlike surd.

Rationalizing Factor: When the product of two surds is rational, then each one of them is called the rationalizing factor of the other.

Unit-2 FACTORIZATION

Linear Polynomial: A polynomial of degree "1" is called a linear polynomial.

Quadratic Polynomial: A polynomial of degree "2" is called a quadratic polynomial.

Cubic Polynomial: A polynomial of degree "3" is called a cubic polynomial.

Types of Factorization: $kx + ky + kz$, $ax + ay + bx + by$, $a^2 \pm 2ab + b^2$

$$a^2 - b^2, (a^2 \pm 2ab + b^2) - c^2, a^4 + a^2b^2 + b^4 \text{ or } a^4 + b^4,$$

$$x^2 + px + q, x^2 + bx + c,$$

$$a^3 + 3a^2bx + 3ab^2 + b^3, a^3 - 3a^2b + 3ab^2 - b^3,$$

$$a^3 \pm b^3.$$

Remainder Theorem: If a polynomial $P(x)$ of degree $n \geq 1$ is divided by a polynomial 'x-a' where 'a' is any constant, then remainder is $P(a)$.

Remainder Theorem: If a polynomial $P(x)$ is divided by 'x-a' such that $P(a) = 0$, then 'x-a' is a factor of $P(x)$.

Unit-3 ALGEBRAIC MANIPULATION

H.C.F: The H.C.F of two or more their two algebraic expressions is the expression of highest degree which divides each of them without remainder.

L.C.M: The least common multiple of two or more algebraic expressions is the expression of lowest degree which is divisible by each of them without remainder.

Unit-4 LINEAR EQUATIONS AND INEQUALITIES

Linear Equation: An equation that can be written in the form $ax + b = 0$, $a \neq 0$ where a and b are constants and x is a variable is called a linear equation in one variable.

Solution of a Linear equation : Any value of the variable, which makes the equation a true statement is called the solution of a linear equation.

Absolute Value: For each real number ' x ' the absolute value of x , denoted by $|x|$, is defined by:

$$|x| = \begin{cases} x, & \text{if } x > 0 \\ 0, & \text{if } x = 0 \\ -x, & \text{if } x < 0 \end{cases}$$

Linear Inequalities: Two algebraic expressions joined by an inequality symbol such as $>$, $<$, $\leq$, $\geq$ is called an inequality.

Trichotomy Property: If $x, y \in R$, then either $x > y$ or $x = y$ or $x < y$.

Transitive Property: If $x, y, z \in R$, then $x > y$ and $y > z \Rightarrow x > z$.

Additive Property: If $\forall a, b, c, d \in R$, then $a > b$ and $c > d \Rightarrow a + b > b + d$ and $c < d$ and $c < d \Rightarrow a + c < b + d$.

Multiplicative Property: $\forall a, b, c, d \in R$, $a > b$ and $c > d \Rightarrow ac > bd$ and $a < d$ and $c > d \Rightarrow ac > bd$.

Unit-5 QUADRATIC EQUATIONS

Quadratic Equation: A quadratic equation in one variable is an equation that can be written in the form $ax^2 + bx + c = 0$, where $a \neq 0$. Here ' x ' is a variable, where a, b and c are real numbers.

Solution of quadratic Equation: We can solve a quadratic equation by
(i) factorization (ii) completing the square method.

Quadratic Formula:
$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Unit-6 MATRICES AND DETERMINANTS

Matrix: A rectangular array of number, enclosed by a pair of brackets and subject to certain rule is called a matrix.

Order of a matrix: The number of rows and columns in a matrix determine its order.

Row matrix: A matrix consisting of one row only is called a row matrix.

Column matrix: A matrix consisting of one column only is called a column matrix.

Square matrix: In a square matrix, the number of rows and columns are equal.

Rectangular matrix: In a rectangular matrix, number of rows and columns are not same.

Zero or null matrix: If all elements in a matrix are zero, the matrix is called a zero or null matrix.

Unit or Identity matrix: In an identity matrix, the diagonal elements are unity and off diagonal elements are all zero.

Transpose of a matrix: A matrix obtained by interchanging rows into columns is called transpose of a matrix.

Symmetric matrix: A matrix A is said to be symmetric, if $A' = A$.

Skew-Symmetric matrix : A matrix A is said to be skew-symmetric, if $A' = -A$.

Determinant: A real number associated with a square matrix is called determinant of a square matrix.

Singular matrix: If the determinant of a square matrix is zero, it is called a singular matrix, other wise non-singular matrix. Adjoint of a square matrix of order 2×2
In the adjoint of a square matrix of order 2×2 , the diagonal elements are interchanged, where as the sign of off diagonal elements are changed.
Multiplicative inverse of a square matrix, A matrix B is said to be multiplications inverse of ' A ', is $AB = I$.

Unit-7 FUNDAMENTALS OF GEOMETRY

Angle: An angle is the union of two rays with common end point.

Right Angle: A right angle contains 90° .

Straight Angle: A straight angle contains 180° .

Acute Angle: An acute angle contains more than 0° and less than 90° .

Obtuse Angle: An obtuse angle contains more than 90° and less than 180° .

Reflex Angle: An reflex angle contains more than 180° and less than 360° .

Equal Angle: Equal angle are angle with equal measures.

Adjacent Angle: Two angles with the common vertex and a common side between them.

Complementary Angle: Two angles whose sum is a right angle.

Supplementary Angle: Two angles whose sum is a straight angle.

Vertical Angle: Two non adjacent angles, each less than a straight angle, formed by two intersecting lines.

Result 1: The sum of the angles of a triangle is a straight angle.

2: If two angles are complements of equal angles, they are equal.

3: If two angles are supplements of the same angle, they are equal.

4: Two lines parallel to a third line are parallel to each other.

5: If three parallel lines in percept congruent segments of one transversal, they intersect congruent segment of every transversal.

6: If a line bisect one side of a triangle and parallel to a second side. It bisect the third side.

Transversal: A transversal is a line that intersects two lines in different points.

Congruent Figures: Two geometrical figures which have the same size and shape are congruent.

Polygon: A polygon is a closed broken line in a plane.

Equilateral Triangle: A triangle with three equal sides.

Isosceles Triangle: A triangle with two equal sides.

Scalene Triangle: A triangle with no equal side.

Right Triangle: A triangle containing one right angle.

Obtuse Triangle: A triangle containing one obtuse angle.

Acute Triangle: A triangle containing three acute angle.

Equiangular Triangle: A triangle containing three equal angle.

Properties for congruency between two Triangle: (i) $SSS \cong SSS$ (ii) $SAS \cong SAS$
(iii) $ASA \cong ASA$ (iv) $AAS \cong AAS$ (v) $RHS \cong RHS$

Quadrilateral: A polygon with four sides.

Parallelogram: A quadrilateral with two pairs of parallel sides.

Rectangle: A parallelogram containing a right angle.

Square: A equilateral rectangle.

Circle: A set of points in a plane which are at a constant distance from a fixed point.

Radius: A segment joining the center to any point on the circle.

Diameter: A chord that passes through the center.

Arc: A portion of a circle consisting of two end points and the set of points on the circle between them.

Semi Circle: An arc which is half of a circle.

Minor Arc: An arc less than a semi-circle.

Major Arc: An arc greater than a semi-circle.

Equal Circles: Circles having equal radii and equal diameters.

Secant Line: A line which intersects a circle in two points.

Tangent: A line perpendicular to the radius of a circle at its outer extremity.

Sector: A circular region bounded by an arc of a circle and its two corresponding radial segments.

Concyclic Points: Points lying on the circumference of the same circle.

Concentric Circles: Circles in the same plane with same center and different radii.

Central Angle: An angle formed at the center of the circle by two radii.

Result: (1) Angle in a semi-circle is a right angle.

(2) Angle in the segment are equal.

(3) All angles inscribed in the same arc are equal in measure.

Unit-8 PRACTICAL GEOMETRY

- 1- An angle bisector of a triangle is a line-segment that bisects an angle of the triangle and has its other end on the side opposite to that angle.
- 2- Every triangle has three angle bisectors, one for each angle.
- 3- An altitude of a triangle is the line-segment from one vertex, perpendicular to the line containing the opposite side.
- 4- Every triangle has three altitudes, one from each vertex.
- 5- A line-segment which bisects any side of a triangle and makes a right angle with the side as its midpoint is called the perpendicular bisector of the side of a triangle.
- 6- Every triangle has three perpendicular bisectors, one for each side.
- 7- The point at which the three angle bisectors of a triangle meet is called the incentre of the triangle.
- 8- The point at which the three altitudes of a triangle meet is called the orthocentre of the triangle.
- 9- The point of intersection of the three perpendicular bisectors of the sides of a triangle is called the circum-centre of the triangle.
- 10- The point at which the three medians of a triangle meet is called the centroid of a triangle.
- 11- A line coplanar with a circle intersecting the circle at one point only is called the tangent line to the circle.

Unit-9 AREAS AND VOLUMES

Pythagoras Theorem: The square of the hypotenuse of a right triangle is equal to the sum of the squares of the legs.

Area: The space inside the boundary of a shape.

Area of a Triangle: $A = \frac{1}{2} \times \text{base} \times \text{altitude}$.

Area of a Triangle: $A = \sqrt{S(s-a)(s-b)(s-c)}$ $S = \frac{a+b+c}{2}$, a, b, c are the sides of a triangle.

Area of an Equilateral Triangle: $A = \frac{\sqrt{3}a^2}{4}$, where 'a' is the side of the triangle.

Area of a Rectangle: $A = \text{length} \times \text{breadth}$.

Area of a Square: $A = \text{side} \times \text{side}$.

Area of a Parallelogram: $A = \text{base} \times \text{altitude}$.

Area of a Circle: $A = \pi r^2$.

Circumference of a Circle: $C = 2\pi r$.

Area of a Semi-Circle: $A = \frac{1}{2}(\pi r^2)$

Area of a Concentric Circle: $A = [r_1^2 - r_2^2]$

r_1 is the radius of outer circle.

r_2 is the radius of inner circle.

Volume: The space/inside/the/boundary/of/a/three/dimensional/shape.

Volume of a Cube: $V = l^3$, l is the length of edge.

Volume of a Cuboid: $V = l \times b \times h$ l = length b = breadth h = height

Volume of a Right Circular Cylinder: $V = \pi r^2 h$

h = height of the cylinder

r = radius of the base

Volume of a Right Circular Cone: $V = \frac{1}{3} \pi r^2 h$

h = height of the cone

r = radius of the base

Volume of Sphere: $V = \frac{4}{3} \pi r^3$

Volume of a Hemisphere: $V = \frac{2}{3} \pi r^3$

Unit-10 INTRODUCTION OF COORDINATE GEOMETRY

Distance Formula: $d = |PQ| = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$

1- A point/in/a/number/plane/determines/a/unique/ordered/pair/of/numbers.

2- With/every/ordered/pair/of/numbers/is/associated/a/unique/point/in/the/place.

Collinear Points: Points/lying/on/the/same/straight/line/are/called/collinear/points.

Non-Collinear Points: Points/which/do/not/lie/on/a/same/straight/line/are/called/non-collinear/points.

SYMBOLS

Symbol	Stands for	Symbol	Stands for
$<$	is/less/than	$\because$	because/as
$>$	is/greater/than	$\therefore$	therefore//so
$\leq$	is/less/than/or/equal/to	$:$	ratio
$\geq$	is/greater/than/or/equal/to	$\therefore$	is/proportional/to
$=$	is/equal/to	∞	varies
$\neq$	is/not/equal/to	$ $	tally/mark
$\nless$	is/not/less/than	Σ	summation
$\ngtr$	is/not/greater/than	$\overline{AB}$	line/segment AB
$\in$	belongs/to	$\overrightarrow{AB}$	ray AB
$\forall$	for/all	$\leftrightarrow$	line
$\sqrt{\quad}$	square/root	$\angle$	angle
$ x $	absolute/value/of x	$\triangle$	triangle
$\Rightarrow$	implies/that	$\sim$	is/similar/to
$\Leftrightarrow$	if/and/only/if	$\cong$	is/congruent/to
$\wedge$	and	$\approx$	is/approximately/equal/to
$\cup$	union	$\parallel$	is/parallel/to
$\vee$	or	$\widehat{AB}$	arc AB
$\subset$	infer/section	$\leftrightarrow$	correspondence

A

Area Of Concentric Circles	261
Absolute Value	97
Acute Angle	177
Add And Subtract Matrices	138
Addition And Subtraction Of Matrices	138
Addition And Subtraction Of Surds	23
Addition Of Matrices	138
Additive Identity Of Matrices	142
Additive Inverse Of A Matrix	143
Adjacent Angles	178
Adjacent, complementary and supplementary angles	178
Adjoint Of A Matrix	157
Algebraic Expressions	2
Algebraic Manipulation	57
Altitudes Of A Triangle	225
Angle	176
Angle Bisectors Of A Triangle	224
Angle In A Semi-circle Is A Right Angle	210
Angle In The Same Segment Are Equal	211
Applications	213
ARC	208
Area Of A Circle	260
Area Of A Parallelogram When Base And Altitude Are Given	255
Area Of A Semicircle	260
Area Of A Triangle When All The Three Sides Are Given	252
Area Of An Equilateral Triangle When Its Side Is Given	254
Area Of Four Walls Of A Room	256
Areas	251
Areas And Volumes	247
Areas Of Rectangular And Square Fields	257
Associative Law	141
Associative Law Of Matrices With Respect To Multiplication	147

B

Basic Operations On The Algebraic Fractions	71
---	----

C

Calculate Unknown Angles	182
Calculate Unknown Angles Of A Triangle	184

Center	206
Central Angle	212
Chord	207
Circle	206
Collinear And Non-collinear Points	281
Collinear Points	281
Collinearity Of Three Points	282
Column Matrix	133
Commutative Law	141
Complementary Angles	179
Concentric Circles	209
Congruent And Similar Figures	193
Congruent Figures	193
Congruent Triangles	198
Conjugate Binomial Surds	27
Construction	222
Construction Of Quadrilaterals	230
Construction Of Triangle	222
Cramer's Rule	167
Cube	264
Cube And Cuboid	264
Cubic Polynomials	36
Cuboid	265

D

Derivation Of Quadratic Formula	115
Determinant Function	156
Diagonal Matrix	134
Diagonals Of A Rectangle Bisect Each Other	204
Diagonals Of A Square Bisect Each Other	203
Diameter	207
Direct Common Tangent Or External Tangent	235
Distance Between Two Points	278
Distance Formula	276
Distributive Laws	149
Division Of A Rational Expression	9
Drawing Tangents	239
Drawing Tangents To Two Equal Circles	235
Drawing Tangents To Two Un-equal Circles	237

E

Equal Angles	178
Equal Circles	208
Equations Involving Absolute Value	97
Equations Involving Radicals	93
Evaluate Determinant Of A Matrix	156
Examine A Given Algebraic Expression	4

F

Factorization	35
Factorization Of Expressions	36
Factorizing A Cubic Polynomial	52
Finding Remainder Without Dividing	49
Formulae	13
Four Angles Of A Rectangle Are Right Angles	204
Four Angles Of A Square Are Right Angles	203
Four Sides Of A Square Are Equal	202
Fundamentals Of Geometry	175

H

H.C.F By Division	61
H.C.F. By Factorization Method	58
Hemispheres	268
Highest Common Factor (HCF) And Least Common Multiple (LCM)	58

I

Improper Rational Expression	3
Inequalities ($>$, $<$) And ($>$, $<$)	98
Introduction To Coordinate Geometry	275
Inverse Of A Non-singular Matrix	160
Irrational Numbers	21

L

L.C.M. By Factorization	64
Laws Of Addition Of Matrices	141
Laws Of Radicals	22
Least Common Multiple (L.C.M)	64
Linear Equation In One Variable	86
Linear Equations	86
Linear Inequalities	98
Linear Polynomials	36

M

Major Arc	208
Matrices And Determinants	127
Matrix Equality	131
Matrix Inversion Method	165
Medians Of A Triangle	228

Minor Arc	208
Mixed Surds	22
Multiplication And Division Of The Algebraic Fractions	72
Multiplication And Division Of Two Surds	23
Multiplication Of Matrices	145
Multiplicative Inverse	158
Multiplicative Inverse Of A Matrix	156

N

Non-singular Matrix	157
---------------------	-----

O

Obtuse Angle	177
Opposite Sides Of A Rectangle Are Equal	203
Order Of A Matrix	130

P

Parallel Lines	187
Parallelogram	231
Perpendicular Bisectors Of The Sides Of A Triangle	227
Problems Involving Quadratic Equations	121
Proper Rational Expression	3
Properties Of A Parallelogram	205
Properties Of Angles	210
Properties Of Congruency	202
Properties Of Congruency Between Two Triangles	198
Properties Of Inequalities	99
Properties Of Parallel Lines	187
Pure Surds	22
Pythagoras Theorem	248

Q

Quadratic Equations	108
Quadratic Polynomials	36
Quadrilaterals	202

R

Radius	206
Rational Expression	2

INDEX

Rational Expression In Its Lowest Terms	4
Rational Numbers	21
Rationalization	27
Rationalizing Factor	27
Rationalizing Of Surds	28
Rationalizing The Denominator	24
Real Numbers	21
Rectangle	230
Rectangular Matrix	133
Reduce A Rational Expression To Its Lowest Terms	5
Reflex Angle	177
Relation Between The Pairs of angles	191
Remainder Theorem And Factor Theorem	47
Right Angle	177
Right Circular Cone	266
Row Matrix	133

S

Scalar Matrix	134
Secant Line	209
Sector	209
Semi Circle	208
Similar Figures	195
Similar Surds	23
Singular And Non-singular Matrices	157
Singular Matrix	157
Skew-symmetric Matrix	136
Solution Of A Linear Equation	87
Solution Of A Quadratic Equation	108
Solution Of A Quadratic Equation By Completing The Square Method	110
Solution Of A Quadratic Equation By Factorization	108
Solution Of Simultaneous Linear Equations	165
Solving Linear Inequalities	101
Square	231
Square Matrix	133
Square Root By Division Method	79
Square Root By Factorization Method	75
Square Root Of Algebraic Expression	75
Straight Angle	176
Subtraction Of Matrices	138
Sum, Difference And Product Of Rational Expressions	6
Supplementary Angles	179

Surds	21
Surds of Radicals	21
Surds Of Second Order	23
Symbol	196
Symmetric Matrix	135

T

Tangent	209
Tangent To A Circle	233
Tangent To Two Un-equal Intersecting Circles	240
Tangent To Two Un-equal Touching Circles	239
The Area Of A Rectangle When Its Two Sides Are Given	255
The Area Of A Triangle	252
The Area Of A Triangle When Base And Altitude Is Given	253
The Factor Theorem	50
The Quadratic Formula	115
The Remainder Theorem	48
The Sphere	267
To Draw Angle-bisectors Of A Triangle	224
Transitive	100
Transpose Of A Matrix	135
Transverse Common Tangent Or Internal Tangent	236
Trichotomy	99
Types Of Matrices	133

U

Unit Matrix Or Identity Matrix	134
Use Of Distance Formula	283
Transverse	189

V

Value Of An Algebraic Expression	10
Vertical Angles	181
Volume	264
Volume Of Right Circular Cylinder	266

Z

Zero Or Null Matrix	134
Zeros Of Polynomial	50

About the Authors


The Author of this book Prof. Dr. Muhammad Ozair Ahmed, Chairman, Department of Mathematics, University of Engineering and Technology (UET) Lahore, has completed his Ph.D from University of Western Ontario, Canada. Beside his 4 years teaching experience in colleges, has 30 years experience of teaching in University. He also taught as teaching Assistant in University of Western Ontario Canada. He has been the President of Punjab Mathematical Society and attended many National and International Seminars on Mathematics.

Before joining UET in 1980 as Lecturer he taught as Lecturer in Govt. College Garhi Dopatta Azad Kashmir and Talimat-e-Alia Girls College Rawalpindi. Then he taught at UET as Assistant Professor, Associate Professor and Professor of Mathematics and Presently he is working as the Chairman Department of Mathematics, UET Lahore.

Mr. Maqsood Raza Ahmad, Associate Professor of Mathematics obtained the Degree of M.Sc in Applied Mathematics from University of the Punjab, Lahore, securing first Position in the Government College, Lahore and was awarded Roll of Honour. In 1979 he joined as lecturer in the Department of Mathematics, University of Balochistan Quetta. There he taught M.Sc classes for a period of almost one year. After having selection by P.P.S.C he appointed in College Cadre in the late 1980. In 1982, he was awarded Postgraduate Scholarship for Bulgaria. In 1990 he was nominated for Commonwealth Scholarship for Postgraduate Studies. In 1994 he was awarded the degree of M.Phil (Fluid Dynamics) by the University of Engineering and Technology, Lahore. The Publications Mr. Maqsood Raza Ahmad has to his credit includes six research papers of postgraduate level in the field of fluid dynamics and more than 20 books of Primary, Middle, Secondary and Intermediate Level. He is presently teaching at Government Shalimar Degree College, Baghbanpura Lahore.



Everybody is sitting in one room whereas the whole house is fully lit.



Publisher's Note

This book is being Published in Accordance with National Curriculum 2006 and National Textbook and Learning Materials Policy 2007 as per International Standards. This book has been Selected by Government of the Punjab as sole Textbook for adoption and distribution in all Government Schools in Punjab.